

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
2 August 2001 (02.08.2001)

PCT

(10) International Publication Number
WO 01/55115 A1(51) International Patent Classification⁷: C07D 213/82,
239/28, 241/28, 213/75, 401/12, 307/68, A61K 31/44,
31/4965, C07D 207/34, A61K 31/341

(21) International Application Number: PCT/US01/02478

(22) International Filing Date: 26 January 2001 (26.01.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/177,648 27 January 2000 (27.01.2000) US(71) Applicant: CYTOVIA, INC. [US/US]; 6650 Nancy
Ridge Drive, San Diego, CA 92121 (US).(72) Inventors: CAI, Sui Xiong; 3623 Berryfield Court, Sand
Diego, CA 92130 (US). DREWE, John, A.; 7069 Murillo
Lane, Carlsbad, CA 92009 (US).(74) Agents: ESMOND, Robert, W. et al.; Sterne, Kessler,
Goldstein & Fox P.L.L.C., 1100 New York Avenue, N.W.,
Suite 600, Washington, DC 20005-3934 (US).(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/55115 A1

(54) Title: SUBSTITUTED NICOTINAMIDES AND ANALOGS AS ACTIVATORS OF CASPASES AND INDUCERS OF APOPTOSIS AND THE USE THEREOF



(V)

(57) Abstract: The present invention is directed to substituted nicotinamides and analogs thereof, represented by Formula (V) or a pharmaceutically acceptable salt or prodrug thereof, wherein: Ar' and Ar are independently optionally substituted aryl or optionally substituted heteroaryl, provided that the ring structure of said optionally substituted heteroaryl comprises not more than two nitrogen atoms; and R₁₁ is hydrogen; or alkyl, cycloalkyl, aryl or heteroaryl, each of which is optionally substituted. The present invention also relates to the

discovery that compounds having Formula (V) are activators of caspases and inducers of apoptosis. Therefore, the compounds of this invention may be used to induce cell death in a variety of clinical conditions in which uncontrolled growth and spread of abnormal cells occurs.

Substituted Nicotinamides and Analogs as Activators of Caspases and Inducers of Apoptosis and the Use Thereof

Field of the Invention

This invention is in the field of medicinal chemistry. In particular, the invention relates to substituted nicotinamides and analogs, and the discovery that these compounds are activators of caspases and inducers of apoptosis. The invention also relates to the use of these compounds as therapeutically effective anti-cancer agents.

Description of Background Art

Organisms eliminate unwanted cells by a process variously known as regulated cell death, programmed cell death or apoptosis. Such cell death occurs as a normal aspect of animal development as well as in tissue homeostasis and aging (Glucksmann, A., *Biol. Rev. Cambridge Philos. Soc.* 26:59-86 (1951); Glucksmann, A., *Archives de Biologie* 76:419-437 (1965); Ellis, *et al.*, *Dev.* 112:591-603 (1991); Vaux, *et al.*, *Cell* 76:777-779 (1994)). Apoptosis regulates cell number, facilitates morphogenesis, removes harmful or otherwise abnormal cells and eliminates cells that have already performed their function. Additionally, apoptosis occurs in response to various physiological stresses, such as hypoxia or ischemia (PCT published application WO96/20721).

There are a number of morphological changes shared by cells experiencing regulated cell death, including plasma and nuclear membrane blebbing, cell shrinkage (condensation of nucleoplasm and cytoplasm), organelle relocation and compaction,

chromatin condensation and production of apoptotic bodies (membrane enclosed particles containing intracellular material) (Orrenius, S., *J. Internal Medicine* 237:529-536 (1995)).

Apoptosis is achieved through an endogenous mechanism of cellular suicide (Wyllie, A.H., in *Cell Death in Biology and Pathology*, Bowen and Lockshin, eds., Chapman and Hall (1981), pp. 9-34). A cell activates its internally encoded suicide program as a result of either internal or external signals. The suicide program is executed through the activation of a carefully regulated genetic program (Wyllie, *et al.*, *Int. Rev. Cyt.* 68:251 (1980); Ellis, *et al.*, *Ann. Rev. Cell Bio.* 7:663 (1991)). Apoptotic cells and bodies are usually recognized and cleared by neighboring cells or macrophages before lysis. Because of this clearance mechanism, inflammation is not induced despite the clearance of great numbers of cells (Orrenius, S., *J. Internal Medicine* 237:529-536 (1995)).

It has been found that a group of proteases are a key element in apoptosis (see, e.g., Thornberry, *Chemistry and Biology* 5:R97-R103 (1998); Thornberry, *British Med. Bull.* 53:478-490 (1996)). Genetic studies in the nematode *Caenorhabditis elegans* revealed that apoptotic cell death involves at least 14 genes, two of which are the pro-apoptotic (death-promoting) *ced* (for *cell death abnormal*) genes, *ced-3* and *ced-4*. CED-3 is homologous to interleukin 1 beta-converting enzyme, a cysteine protease, which is now called caspase-1. When these data were ultimately applied to mammals, and upon further extensive investigation, it was found that the mammalian apoptosis system appears to involve a cascade of caspases, or a system that behaves like a cascade of caspases. At present, the caspase family of cysteine proteases comprises 14 different members, and more may be discovered in the future. All known caspases are synthesized as zymogens that require cleavage at an aspartyl residue prior to forming the active enzyme. Thus, caspases are capable of activating other caspases, in the manner of an amplifying cascade.


Apoptosis and caspases are thought to be crucial in the development of cancer (*Apoptosis and Cancer Chemotherapy*, Hickman and Dive, eds., Humana Press (1999)). There is mounting evidence that cancer cells, while containing caspases, lack parts of the molecular machinery that activates the caspase cascade. This makes the cancer cells lose their capacity to undergo cellular suicide and the cells become cancerous. In the case of the apoptosis process, control points are known to exist that represent points for

intervention leading to activation. These control points include the CED-9-BCL-like and CED-3-ICE-like gene family products, which are intrinsic proteins regulating the decision of a cell to survive or die and executing part of the cell death process itself, respectively (see, Schmitt, *et al.*, *Biochem. Cell. Biol.* 75:301-314 (1997)). BCL-like proteins include BCL-xL and BAX-alpha, which appear to function upstream of caspase activation. BCL-xL appears to prevent activation of the apoptotic protease cascade, whereas BAX-alpha accelerates activation of the apoptotic protease cascade.

It has been shown that chemotherapeutic (anti-cancer) drugs can trigger cancer cells to undergo suicide by activating the dormant caspase cascade. This may be a crucial aspect of the mode of action of most, if not all, known anticancer drugs (Los, *et al.*, *Blood* 90:3118-3129 (1997); Friesen, *et al.*, *Nat. Med.* 2:574 (1996)). The mechanism of action of current antineoplastic drugs frequently involves an attack at specific phases of the cell cycle. In brief, the cell cycle refers to the stages through which cells normally progress during their lifetimes. Normally, cells exist in a resting phase termed G₀. During multiplication, cells progress to a stage in which DNA synthesis occurs, termed S. Later, cell division, or mitosis occurs, in a phase called M. Antineoplastic drugs such as cytosine arabinoside, hydroxyurea, 6-mercaptopurine, and methotrexate are S phase specific, whereas antineoplastic drugs such as vincristine, vinblastine, and paclitaxel are M phase specific. Many slow growing tumors, for example colon cancers, exist primarily in the G₀ phase, whereas rapidly proliferating normal tissues, for example bone marrow, exist primarily in the S or M phase. Thus, a drug like 6-mercaptopurine can cause bone marrow toxicity while remaining ineffective for a slow growing tumor. Further aspects of the chemotherapy of neoplastic diseases are known to those skilled in the art (See, e.g., Hardman, *et al.*, eds., *Goodman and Gilman's The Pharmacological Basis of Therapeutics*, Ninth Edition, McGraw-Hill, New York (1996), pp. 1225-1287). Thus, it is clear that the possibility exists for the activation of the caspase cascade, although the exact mechanisms for doing so are not clear at this point. It is equally clear that insufficient activity of the caspase cascade and consequent apoptotic events are implicated in various types of cancer. The development of caspase cascade activators and inducers of apoptosis is a highly desirable goal in the development of therapeutically effective antineoplastic agents. Moreover, since autoimmune disease and certain degenerative diseases also

involve the proliferation of abnormal cells, therapeutic treatment for these diseases could also involve the enhancement of the apoptotic process through the administration of appropriate caspase cascade activators and inducers of apoptosis.

PCT published patent application WO95/25723 discloses anilide derivatives as fungicides:



wherein,

X is O or S;

A is a 6 membered heteroaryl group comprising at least one nitrogen atom, which is optionally substituted by one or more of the group R²;

R¹ is alkyl, cycloalkyl, cycloalkenyl, alkenyl, alkynyl, or amino, (each of which is optionally substituted), Y¹-X-, halogen, cyano, nitro, acyl, acyloxy, optionally substituted heterocyclyl or optionally substituted phenyl; or two adjacent groups together with the carbon atoms to which they are attached can form an optionally substituted benzo ring;

R² is the same meaning as R¹ or two adjacent groups together with the carbon atoms to which they are attached can form an optionally substituted heterocyclic ring;

Y is alkyl, cycloalkyl, cycloalkenyl, alkenyl or alkynyl, each of which is optionally substituted, hydrogen or acyl;

Y¹ has the same meaning as Y or is optionally substituted phenyl or optionally substituted heterocyclyl;

Z is C(=X¹)-X²-R³, cyano, nitro, amino, acyl, optionally substituted heterocyclyl, -C(R⁵)=N-OR⁶ or -C(R⁵)=N-NR⁶R⁷;

R³ is alkyl, cycloalkyl, cycloalkenyl, alkenyl, alkynyl, phenyl or heterocyclyl, each of which is optionally substituted, hydrogen or an inorganic or organic cationic group;

-5-

X^1 and X^2 , which may be the same or different, are O or S;

R^5 , R^6 and R^7 , which may be the same or different, are alkyl, cycloalkyl, cycloalkenyl, alkenyl, alkynyl, phenyl or heterocyclyl, each of which is optionally substituted or hydrogen or R^6 and R^7 together with the atom(s) to which they are attached can form a ring;


and n is 0 to 4,

together with complexes with metal salts, as well as salts with bases of compounds which are acids and salts with acids of compounds which are bases, with the proviso that when Y is hydrogen and

i) when Z is carboxy, methoxycarbonyl or ethoxycarbonyl ring A is not unsubstituted pyridyl or pyrazinyl; and

ii) when Z is carboxy and n is 0, A is not 2-chloro-3-pyridyl, 6-(2-diethylaminoethoxy)-3-pyridyl or a 2-pyridyl group.

PCT WO9936391 discloses benzenesulfonamide, benzamide, diarylsulfone and benzophenone compounds as pharmacological agents in the treatment of cancer, psoriasis, vascular restenosis, infections, atherosclerosis and hypercholesterolemia:



wherein, A represents N or C- R^1 , B represents N or C- R^5 ; and

R^1 and R^5 independently represent hydrogen, halogen, (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl, -OR⁶, -NR⁶R⁷, -S(O)_mR⁶, -CN, -NO₂, -S(O)_nNR⁶R⁷, or -N₃; wherein R^6 and R^7 are independently selected from hydrogen, (C₁-C₈)alkyl, (C₁-C₈)heteroalkyl;

m is an integer of from 0 to 3;

n is an integer of from 1 to 2;

R^2 and R^3 are independently -OR⁸, -SR⁸, and -NR⁸R⁹, wherein R^8 and R^9 are independently hydrogen, (C₁-C₆)alkyl, (C₁-C₆)heteroalkyl;

-6-

R^4 is hydrogen, (C_1-C_8) alkyl, (C_1-C_8) heteroalkyl, $-OR^{10}$, $-SR^{10}$, or $-NR^{10}R^{11}$; wherein R^{10} and R^{11} are independently hydrogen, (C_1-C_6) alkyl, (C_1-C_6) heteroalkyl;

optionally, R^1 is linked to R^2 to form a fused ring, R^2 is linked to R^3 to form a fused ring, or R^2 is linked to both R^1 and R^3 to form two additional fused rings;


X represents $-S(O)_p$ or $-C(O)-$, wherein p is 1 or 2;

Y represents a single bond, $-CH_2-$ or $-N(R^{12})-$, wherein R^{12} is selected from hydrogen, (C_1-C_6) alkyl, (C_1-C_6) heteroalkyl and arylalkyl; and

Z represents an aryl group or an arylalkyl group.

Summary of the Invention

The present invention is related to the discovery that substituted nicotinamides and nicotinamide analogs as represented in Formula V are activators of the caspase cascade and inducers of apoptosis:



or a pharmaceutically acceptable salt or prodrug thereof, wherein:

Ar' and Ar are independently optionally substituted aryl or optionally substituted heteroaryl, provided that the ring structure of said optionally substituted heteroaryl comprises not more than two nitrogen atoms; and

R_{11} is hydrogen; or alkyl, cycloalkyl, aryl or heteroaryl, each of which is optionally substituted.

Thus, an aspect of the present invention is directed to the use of compounds of Formula V as inducers of apoptosis.

A second aspect of the present invention is to provide a method for treating, preventing or ameliorating neoplasia and cancer by administering a compound of Formula V to a mammal in need of such treatment.

Many of the compounds within the scope of the present invention are novel compounds. Therefore, a third aspect of the present invention is to provide novel compounds of Formula V, and to also provide for the use of these novel compounds for treating, preventing or ameliorating neoplasia and cancer.

5 A fourth aspect of the present invention is to provide a pharmaceutical composition useful for treating disorders responsive to the induction of apoptosis, containing an effective amount of a compound of Formula V in admixture with one or more pharmaceutically acceptable carriers or diluents.

10 A fifth aspect of the present invention is directed to methods for the preparation of novel compounds of Formula V.

Brief Description of the Drawings

15 Figs. 1A-B depict fluorescent micrographs of Jurkat cells as controls and as treated with drug and stained with a fluorescent DNA probe, Syto16. Fig. 1A depicts control cells. Fig. 1B depicts cells treated with 5 μ M of N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide for 24 h, showing shrunken and fragmented nuclei.

20 Fig. 2 is a graph showing mitotic arrest in Jurkat cells treated for 6 h with different concentrations of N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide. Fig. 2 shows increasing percent of mitotic arrest with increasing drug concentration up to a concentration of 5 μ M.

25 Figs. 3A-B are graphs showing drug induced cell cycle arrest and apoptosis in T47D cells. Fig. 3A: control cells showing most of the cells in G1(M2). Fig. 3B: cells treated with 10 μ M of N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide for 48 h showing a reduction in the G1(M2), an increase in the G2/M (M4) and sub-diploid DNA content of cells (M1).

30 Figs. 4A-B are graphs showing drug induced cell cycle arrest and apoptosis in T47D cells. Fig. 4A: control cells showing most of the cells in G1(M2). Fig. 4B: cells treated with 1 μ M of 6-chloro-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide for 48 h showing a reduction in the G1(M2), an increase in the G2/M (M4) and sub-diploid DNA content of cells (M1).


Fig. 5 is a graph showing inhibition of proliferation of HeLa cells treated for 48 h with different concentrations of 6-chloro-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide. Fig. 5 shows that 6-chloro-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide inhibit colony formation of HeLa cells with an IC₅₀ of about 100 nM.

5

Detailed Description of the Invention

The present invention arises out of the discovery that substituted nicotinamides and nicotinamide analogs, as represented in Formula V, are potent and highly efficacious activators of the caspase cascade and inducers of apoptosis:

10



or a pharmaceutically acceptable salt or prodrug thereof, wherein:

15


Ar' and Ar are independently optionally substituted aryl or optionally substituted heteroaryl, provided that the ring structure of said optionally substituted heteroaryl comprises not more than two nitrogen atoms; and

R₁₁ is hydrogen; or alkyl, cycloalkyl, aryl or heteroaryl, each of which is optionally substituted.

20

Therefore compounds of Formula V are useful for treating disorders responsive to induction of apoptosis.

Specifically, compounds useful in this aspect of the present invention are represented by Formula I:



25

or pharmaceutically acceptable salts or prodrugs thereof, wherein:

A is N or C-R₈; B is N or C-R₉, D is N or C-R₁₀, E is N or C-R₆, F is N or C-R₇, provided that not more than two of A, B, D, E, and F are N at the same time;

Ar is optionally substituted and is aryl or heteroaryl;

5 R₆-R₁₀ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, aminoalkyl, cyano, cyanoalkyl, acyl, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, alkoxycarbonyl, aryloxy, arylalkoxy, haloalkoxy, carboxy, carbonylamido or alkylthiol; and


10 R₁₁ is hydrogen; or alkyl, cycloalkyl, aryl or heteroaryl, each of which is optionally substituted.

Preferred compounds of Formula I include compounds wherein Ar is optionally substituted phenyl, naphthyl, pyridyl, quinolyl, isoquinolyl, thienyl, furyl, or pyrrolyl.

15 Preferred compounds of Formula I also include compounds wherein A is N, B is C-R₉, D is C-R₁₀, E is C-R₆ and F is C-R₇. Preferred compounds of Formula I also include compounds wherein A and D are N, B is C-R₉, E is C-R₆ and F is C-R₇. Preferred compounds of Formula I also include compounds wherein A and E are N, B is C-R₉, D is C-R₁₀ and F is C-R₇. Preferred compounds of Formula I also include compounds wherein

20 R₁₁ is hydrogen.


Preferred structures of Formula I are substituted nicotinamides and analogs represented by Formulae II-IV. In particular, a preferred embodiment is represented by Formula II:



25 or pharmaceutically acceptable salts or prodrugs thereof, wherein R₇, R₉, R₁₁, D, E and Ar are as defined previously with respect to Formula I.

Preferred compounds falling within the scope of Formula II include compounds wherein Ar is optionally substituted phenyl or pyridyl. Preferred compounds of Formula II also include compounds wherein D is C-R₁₀ and E is C-R₆. Preferred compounds of Formula II also include compounds wherein D are N and E is C-R₆. Preferred compounds of Formula II also include compounds wherein E are N and D is C-R₁₀. Preferred compounds of Formula II also include compounds wherein R₆-R₇ and R₁₀ are independently hydrogen or fluoro.

Another preferred embodiment is represented by Formula III:



or pharmaceutically acceptable salts or prodrugs thereof, wherein R₇, R₉, R₁₁, D, and E are as defined previously with respect to Formula I;


R₁-R₅ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, aminoalkyl, cyano, cyanoalkyl, acyl, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, alkoxycarbonyl, aryloxy, arylalkoxy, haloalkoxy, carboxy, carbonylamido or alkylthiol; or

R₁ and R₂, or R₂ and R₃, or R₃ and R₄, or R₄ and R₅ may be taken together to form a carbocycle or heterocycle, including -OCH₂O-, -(CH₂)₃-, -(CH₂)₄-, -OCH₂CH₂O-, -CH₂N(R)CH₂-, -CH₂CH₂N(R)CH₂-, -CH₂N(R)CH₂CH₂-, -CH=CH-CH=CH-, -N(R)-CH=CH-, -CH=CH-N(R)-, -O-CH=CH-, -CH=CH-O-, -S-CH=CH-, -CH=CH-S-, -N=CH-CH=CH-, -CH=N-CH=CH-, -CH=CH-N=CH-, -CH=CH-CH=N- and -N=CH-CH=N-, wherein the carbocycle or heterocycle is optionally substituted, and R is hydrogen, C₁₋₁₀ alkyl, haloalkyl, aryl, fused aryl,

carbocyclic, a heterocyclic group, a heteroaryl group, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl or aminoalkyl.

Preferred compounds falling within the scope of Formula III include compounds wherein D is C-R₁₀ and E is C-R₆. Preferred compounds of Formula II also include compounds wherein D are N and E is C-R₆. Preferred compounds of Formula II also include compounds wherein E are N and D is C-R₁₀. Preferred compounds of Formula II also include compounds wherein R₆-R₇ and R₁₀ are hydrogen or fluoro. Preferred compounds of Formula III also include compounds wherein R₁ and R₃ is not hydrogen. Preferred compounds of Formula III also include compounds wherein R₂ and R₄ are hydrogen or fluoro. Another group of preferred compounds of Formula III include compounds wherein R₁₁ is hydrogen.

Another preferred embodiment is represented by Formula IV:




or pharmaceutically acceptable salts or prodrugs thereof, wherein R₁, R₃, R₅, R₉, D and E are as defined previously with respect to Formulae I and III.

Preferred compounds falling within the scope of Formula IV include compounds wherein D is C-R₁₀ and E is C-R₆. Preferred compounds of Formula II also include compounds wherein D are N and E is C-R₆. Preferred compounds of Formula II also include compounds wherein E are N and D is C-R₁₀. Preferred compounds of Formula IV also include compounds wherein R₁ and R₃ is not hydrogen. Especially preferred compounds of Formula IV include compounds wherein R₁ is nitro, cyano, trifluoromethyl and methyl. Especially preferred compounds of Formula IV also include compounds wherein R₃ is halo, haloalkyl, alkyl, amino, cyano, acyloxy, azido, alkoxy, aryloxy, arylalkoxy, haloalkoxy, or alkylthiol.

-12-

Another preferred embodiment is represented by Formula VI:




or a pharmaceutically acceptable salt or prodrug thereof, wherein

R₁-R₅, R₇ and R₉-R₁₀ are independently hydrogen, halo, haloalkyl, haloalkoxy, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, aminoalkyl, cyano, cyanoalkyl, acyl, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, alkoxycarbonyl, aryloxy, arylalkoxy, carboxy, carbonylamido or alkylthiol; and

R₁₁ is hydrogen; or alkyl, cycloalkyl, aryl or heteroaryl, each of which is optionally substituted.

Another preferred embodiment is represented by Formula VII:



or a pharmaceutically acceptable salt or prodrug thereof, wherein:

R₁-R₃, R₅-R₁₀ are independently hydrogen, halo, haloalkyl, haloalkoxy, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, aminoalkyl, cyano,

cyanoalkyl, acyl, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, alkoxycarbonyl, aryloxy, arylalkoxy, carboxy, carbonylamido or alkylthiol; and R₁₁ is hydrogen; or alkyl, cycloalkyl, aryl or heteroaryl, each of which is optionally substituted.


5 Exemplary preferred compounds that may be employed in the method of the invention include, without limitation:

N-(4-Methoxy-2-nitrophenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
10 6-Chloro-*N*-(4-methyl-2-nitrophenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(4-methoxy-2-nitrophenyl)-1-*N*-oxide-3-pyridinecarboxamide;
6-Chloro-*N*-(4-chloro-2-nitrophenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(4,5-difluoro-2-nitrophenyl)-3-pyridinecarboxamide;
6-Fluoro-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
15 6-Chloro-*N*-(4-fluoro-2-nitrophenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(2-nitro-4-trifluoromethylphenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(3-bromo-4-methoxy-6-nitrophenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(2-nitro-4-trifluoromethoxyphenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(4-benzyloxy-2-nitrophenyl)-3-pyridinecarboxamide;
20 6-Methyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
5,6-Dichloro-*N*-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(2-methyl-4-methoxyphenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(4-cyano-2-nitrophenyl)-3-pyridinecarboxamide;
4-Chloro-*N*-(4-ethoxy-2-nitrophenyl)-benzoylamide;
25 6-Chloro-*N*-(4-ethoxy-2-nitrophenyl)-*N*-methyl-3-pyridinecarboxamide;
6-Chloro-*N*-(2-cyano-4,5-dimethoxyphenyl)-3-pyridinecarboxamide;
N-(4-Ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
N-(4-Methyl-2-nitrophenyl)-3-pyridinecarboxamide;
6-(2,2,2-Trifluoroethoxy)-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
30 *N*-(4-Ethoxy-2-nitrophenyl)-2-pyridinecarboxamide;
N-(4-Ethoxy-2-nitrophenyl)-5-pyrimidinecarboxamide;

6-Dimethylamino-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(4-*t*-butyl-2-nitrophenyl)-3-pyridinecarboxamide;
N-(4-Ethoxy-2-nitrophenyl)-1-*N*-oxide-2-pyridinecarboxamide;
6-Trifluoromethyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
5 6-Chloro-*N*-(4-chloro-2-trifluoromethylphenyl)-3-pyridinecarboxamide;
6-Methyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyrazinecarboxamide;
6-Chloro-*N*-(4-chloro-2-cyanophenyl)-3-pyridinecarboxamide;
N-(4-Ethoxy-2-nitrophenyl)-3-pyrazinecarboxamide;
6-Chloro-*N*-(2,4-dimethyl-6-nitrophenyl)-3-pyridinecarboxamide;
10 4-Chloromethyl-*N*-(4-ethoxy-2-nitrophenyl)-benzoylamide;
6-Chloro-*N*-(3,4-dimethoxy-6-nitrophenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(2-pyrazinyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(1-*N*-oxide-3-cyano-5-chloromethyl-2-pyrazinyl)-3-
pyridinecarboxamide;
15 6-Chloro-*N*-(2-cyano-4-methylphenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(4-chloro-2-methyl-6-nitrophenyl)-3-pyridinecarboxamide;
4-Trifluoromethyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
6-Chloromethyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
4-Chloro-2-nitro-*N*-(6-chloro-3-pyridyl)-benzoylamide;
20 4-Chloro-2-nitro-*N*-(6-methoxy-3-pyridyl)-benzoylamide;
4-Bromomethyl-3-nitro-*N*-(6-chloro-3-pyridyl)-benzoylamide;
N-(4-Ethoxy-2-nitrophenyl)-3-furancarboxamide; and
N-(4-Ethoxy-2-nitrophenyl)-3-pyrrolecarboxamide.

The present invention is also directed to novel compounds within the scope of
25 Formulae I-VII. In one preferred embodiment, the novel compounds of the present
invention are compounds of Formula III:

-15-



or pharmaceutically acceptable salts or prodrugs thereof, wherein:

R_1 and R_5 are independently selected from the group consisting of hydrogen, hydroxy, alkyl, alkoxy, halogen, NO_2 , cyano, haloalkyl, haloalkoxy, amino and aminoalkyl, provided that at least one of R_1 and R_5 is selected from the group consisting of NO_2 , cyano, alkyl and haloalkyl;

R_2 and R_4 are independently selected from the group consisting of hydrogen, hydroxy, halogen, cyano, haloalkyl, haloalkoxy, amino and aminoalkyl;

R_3 is alkyl, Cl, F, haloalkyl, alkoxy, arylalkoxy, cyano, haloalkyloxy, amino or aminoalkyl;

R_6 is hydrogen, hydroxy, alkyl, NO_2 , cyano, haloalkyl, haloalkyloxy, amino or aminoalkyl;


R_7 is hydrogen, hydroxy, alkyl, NO_2 , cyano, haloalkyl, haloalkyloxy, amino or aminoalkyl;

R_9 is hydroxy, alkyl, halogen, NO_2 , haloalkyl, alkoxy, cyano, haloalkyloxy, amino or aminoalkyl;

R_{10} is hydrogen, hydroxy, alkyl, Cl, F, NO_2 , cyano, haloalkyl, haloalkyloxy, amino or aminoalkyl; and

R_{11} is hydrogen, alkyl or haloalkyl.

In a further preferred embodiment, the compounds of the present invention are compounds of Formula IV:




or pharmaceutically acceptable salts or prodrugs thereof, wherein:

R_3 is alkyl, Cl, F, haloalkyl, alkoxy, arylalkoxy, cyano, haloalkyloxy, amino or aminoalkyl; and

5 R_9 is hydroxy, alkyl, halogen, NO_2 , haloalkyl, alkoxy, cyano, haloalkyloxy, amino or aminoalkyl.

In another preferred embodiment, the compounds of the present invention are compounds of Formula (VI):



10 or pharmaceutically acceptable salts or prodrugs thereof, wherein


R_1 - R_5 , R_7 and R_9 - R_{10} are independently hydrogen, halo, haloalkyl, haloalkoxy, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, aminoalkyl, cyano, cyanoalkyl, acyl, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, alkoxycarbonyl, aryloxy, arylalkoxy, carboxy, carbonylamido or alkylthiol; and

15 R_{11} is hydrogen; or alkyl, cycloalkyl, aryl or heteroaryl, each of which is optionally substituted.

In another preferred embodiment, the novel compounds of the present invention are compounds of Formula (VII):

20

-17-



or pharmaceutically acceptable salts or prodrugs thereof, wherein:

R_1 - R_3 , R_5 - R_{10} are independently hydrogen, halo, haloalkyl, haloalkoxy, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, aminoalkyl, cyano, cyanoalkyl, acyl, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, alkoxycarbonyl, aryloxy, arylalkoxy, carboxy, carbonylamido or alkylthiol; and

R_{11} is hydrogen; or alkyl, cycloalkyl, aryl or heteroaryl, each of which is optionally substituted.

More preferably at least one of R_6 and R_7 is selected from the group consisting of NO_2 , cyano, alkyl and haloalkyl.

Exemplary preferred compounds of the present invention include, without limitation:

- 6-Chloro-*N*-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide;
- 6-Chloro-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
- 6-Chloro-*N*-(4-methyl-2-nitrophenyl)-3-pyridinecarboxamide;
- 6-Chloro-*N*-(4-methoxy-2-nitrophenyl)-1-*N*-oxide-3-pyridinecarboxamide;
- 6-Chloro-*N*-(4-chloro-2-nitrophenyl)-3-pyridinecarboxamide;
- 6-Chloro-*N*-(4,5-difluoro-2-nitrophenyl)-3-pyridinecarboxamide;
- 6-Fluoro-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
- 6-Chloro-*N*-(4-fluoro-2-nitrophenyl)-3-pyridinecarboxamide;
- 6-Chloro-*N*-(4-trifluoromethyl-2-nitrophenyl)-3-pyridinecarboxamide;
- 6-Chloro-*N*-(3-bromo-4-methoxy-6-nitrophenyl)-3-pyridinecarboxamide;
- 6-Chloro-*N*-(2-nitro-4-trifluoromethoxyphenyl)-3-pyridinecarboxamide;
- 6-Chloro-*N*-(4-benzyloxy-2-nitrophenyl)-3-pyridinecarboxamide;

- 6-Methyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
5,6-Dichloro-*N*-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(2-methyl-4-methoxyphenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(4-cyano-2-nitrophenyl)-3-pyridinecarboxamide;
5 4-Chloro-*N*-(4-ethoxy-2-nitrophenyl)-benzoylamide;
6-Chloro-*N*-(4-ethoxy-2-nitrophenyl)-*N*-methyl-3-pyridinecarboxamide;
6-Chloro-*N*-(2-cyano-4,5-dimethoxyphenyl)-3-pyridinecarboxamide;
6-(2,2,2-Trifluoroethoxy)-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
N-(4-Ethoxy-2-nitrophenyl)-2-pyridinecarboxamide;
10 *N*-(4-Ethoxy-2-nitrophenyl)-5-pyrimidinecarboxamide;
6-Dimethylamino-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(4-*t*-butyl-2-nitrophenyl)-3-pyridinecarboxamide;
N-(4-Ethoxy-2-nitrophenyl)-1-*N*-oxide-2-pyridinecarboxamide;
6-Trifluoromethyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
15 6-Chloro-*N*-(4-chloro-2-trifluoromethylphenyl)-3-pyridinecarboxamide;
6-Methyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyrazinecarboxamide;
6-Chloro-*N*-(4-chloro-2-cyanophenyl)-3-pyridinecarboxamide;
N-(4-Ethoxy-2-nitrophenyl)-3-pyrazinecarboxamide;
6-Chloro-*N*-(2,4-dimethyl-6-nitrophenyl)-3-pyridinecarboxamide;
20 4-Chloromethyl-*N*-(4-ethoxy-2-nitrophenyl)-benzoylamide;
6-Chloro-*N*-(3,4-dimethoxy-6-nitrophenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(2-pyrazinyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(1-*N*-oxide-3-cyano-5-chloromethyl-2-pyrazinyl)-3-
pyridinecarboxamide;
25 6-Chloro-*N*-(2-cyano-4-methylphenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(4-chloro-2-methyl-6-nitrophenyl)-3-pyridinecarboxamide;
4-Trifluoromethyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
6-Chloromethyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
4-Chloro-2-nitro-*N*-(6-chloro-3-pyridyl)-benzoylamide;
30 4-Chloro-2-nitro-*N*-(6-methoxy-3-pyridyl)-benzoylamide; and
4-Bromomethyl-3-nitro-*N*-(6-chloro-3-pyridyl)-benzoylamide.

Useful alkyl groups include straight-chained and branched C₁₋₁₀ alkyl groups, more preferably C₁₋₆ alkyl groups. Typical C₁₋₁₀ alkyl groups include methyl, ethyl, propyl, isopropyl, butyl, *sec*-butyl, *tert*-butyl, 3-pentyl, hexyl and octyl groups, which may be optionally substituted.

5 Useful alkoxy groups include oxygen substituted by one of the C₁₋₁₀ alkyl groups mentioned above, which may be optionally substituted.

Useful alkylthio groups include sulphur substituted by one of the C₁₋₁₀ alkyl groups mentioned above, which may be optionally substituted. Also included are the sulfoxides and sulfones of such alkylthio groups.

10 Useful amino groups include -NH₂, -NHR₁₅ and -NR₁₅R₁₆, wherein R₁₅ and R₁₆ are C₁₋₁₀ alkyl or cycloalkyl groups, or R₁₅ and R₁₆ are combined with the N to form a ring structure, such as a piperidine, or R₁₅ and R₁₆ are combined with the N and other group to form a ring, such as a piperazine. The alkyl group may be optionally substituted.

15 Optional substituents on the alkyl groups include one or more halo, hydroxy, carboxyl, amino, nitro, cyano, C₁-C₆ acylamino, C₁-C₆ acyloxy, C₁-C₆ alkoxy, aryloxy, alkylthio, C₆-C₁₀ aryl, C₄-C₇ cycloalkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₆-C₁₀ aryl(C₂-C₆)alkenyl, C₆-C₁₀ aryl(C₂-C₆)alkynyl, saturated and unsaturated heterocyclic or heteroaryl.

20 Optional substituents on the aryl, aralkyl and heteroaryl groups include one or more halo, C₁-C₆ haloalkyl, C₆-C₁₀ aryl, C₄-C₇ cycloalkyl, C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₆-C₁₀ aryl(C₁-C₆)alkyl, C₆-C₁₀ aryl(C₂-C₆)alkenyl, C₆-C₁₀ aryl(C₂-C₆)alkynyl, C₁-C₆ hydroxyalkyl, nitro, amino, ureido, cyano, C₁-C₆ acylamino, hydroxy, thiol, C₁-C₆ acyloxy, azido, C₁-C₆ alkoxy or carboxy.

25 The term "aryl" as employed herein by itself or as part of another group refers to monocyclic, bicyclic or tricyclic aromatic groups containing from 6 to 14 carbons in the ring portion.

Useful aryl groups include C₆₋₁₄ aryl, preferably C₆₋₁₀ aryl. Typical C₆₋₁₄ aryl groups include phenyl, naphthyl, phenanthrenyl, anthracenyl, indenyl, azulenyl, biphenyl, biphenylenyl and fluorenyl groups.

30 Useful cycloalkyl groups are C₃₋₈ cycloalkyl. Typical cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl.

Useful saturated or partially saturated carbocyclic groups are cycloalkyl groups as described above, as well as cycloalkenyl groups, such as cyclopentenyl, cycloheptenyl and cyclooctenyl.

Useful halo or halogen groups include fluorine, chlorine, bromine and iodine.

5 Useful arylalkyl groups include any of the above-mentioned C₁₋₁₀ alkyl groups substituted by any of the above-mentioned C₆₋₁₄ aryl groups. Preferably the arylalkyl group is benzyl, phenethyl or naphthylmethyl.

10 Useful haloalkyl groups include C₁₋₁₀ alkyl groups substituted by one or more fluorine, chlorine, bromine or iodine atoms, e.g., fluoromethyl, difluoromethyl, trifluoromethyl, pentafluoroethyl, 1,1-difluoroethyl, chloromethyl, chlorofluoromethyl and trichloromethyl groups.

15 Useful acylamino (acylamido) groups are any C₁₋₆ acyl (alkanoyl) attached to an amino nitrogen, e.g., acetamido, chloroacetamido, propionamido, butanoylamido, pentanoylamido and hexanoylamido, as well as aryl-substituted C₁₋₆ acylamino groups, e.g., benzoylamido, and pentafluorobenzoylamido.

Useful acyloxy groups are any C₁₋₆ acyl (alkanoyl) attached to an oxy (—O—) group, e.g., formyloxy, acetoxy, propionyloxy, butanoyloxy, pentanoyloxy and hexanoyloxy.

20 The term heterocycle is used herein to mean a saturated or partially saturated 3-7 membered monocyclic, or 7-10 membered bicyclic ring system, which consists of carbon atoms and from one to four heteroatoms independently selected from the group consisting of O, N, and S, wherein the nitrogen and sulfur heteroatoms can be optionally oxidized, the nitrogen can be optionally quaternized, and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring, and wherein the heterocyclic ring can be substituted on carbon or on a nitrogen atom if the resulting
25 compound is stable.

Useful saturated or partially saturated heterocyclic groups include tetrahydrofuranyl, pyranal, piperidinyl, piperazinyl, pyrrolidinyl, imidazolidinyl, imidazoliny, indoliny, isoindoliny, quinuclidiny, morpholiny, isochromanyl, chromanyl, pyrazolidiny, pyrazoliny, tetronoyl and tetramoyl groups.

The term "heteroaryl" as employed herein refers to groups having 5 to 14 ring atoms; 6, 10 or 14 π electrons shared in a cyclic array; and containing carbon atoms and 1, 2 or 3 oxygen, nitrogen or sulfur heteroatoms.

Useful heteroaryl groups include thienyl, benzo[b]thienyl, naphtho[2,3-b]thienyl, thianthrenyl, furyl, pyranyl, isobenzofuranyl, chromenyl, xanthenyl, phenoxanthiyl, 2H-pyrrolyl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indoliziny, isoindolyl, 3H-indolyl, indolyl, indazolyl, purinyl, 4H-quinoliziny, isoquinolyl, quinolyl, phthalziny, naphthyridinyl, quinozaliny, cinnoliny, pteridinyl, carbazolyl, β -carbolinyl, phenanthridinyl, acridinyl, perimidinyl, phenanthrolinyl, phenazinyl, isothiazolyl, phenothiazinyl, isoxazolyl, furazanyl, phenoxazinyl, 1,4-dihydroquinoxaline-2,3-dione, 7-aminoisocoumarin, pyrido[1,2-a]pyrimidin-4-one, 1,2-benzisoxazol-3-yl, benzimidazolyl, 2-oxindolyl and 2-oxobenzimidazolyl. Where the heteroaryl group contains a nitrogen atom in a ring, such nitrogen atom may be in the form of an N-oxide, e.g., a pyridyl N-oxide, pyrazinyl N-oxide and pyrimidinyl N-oxide.

Some of the compounds of the present invention may exist as stereoisomers including optical isomers. The invention includes all stereoisomers and both the racemic mixtures of such stereoisomers as well as the individual enantiomers that may be separated according to methods that are well known to those of ordinary skill in the art.


Examples of pharmaceutically acceptable addition salts include inorganic and organic acid addition salts such as hydrochloride, hydrobromide, phosphate, sulphate, citrate, lactate, tartrate, maleate, fumarate, mandelate and oxalate; and inorganic and organic base addition salts with bases such as sodium hydroxy, Tris(hydroxymethyl)aminomethane (TRIS, tromethane) and N-methyl-glucamine.

Examples of prodrugs of the compounds of the invention include the simple esters of carboxylic acid containing compounds (e.g., those obtained by condensation with a C_{1-4} alcohol according to methods known in the art); esters of hydroxy containing compounds (e.g., those obtained by condensation with a C_{1-4} carboxylic acid, C_{3-6} dioic acid or anhydride thereof such as succinic and fumaric anhydrides according to methods known in the art); imines of amino containing compounds (e.g., those obtained by condensation with a C_{1-4} aldehyde or ketone according to methods known in the art); carbamate of amino containing compounds such as those described by Leu, *et. al.*, (*J. Med. Chem.* 42:3623-

3628 (1999)) and Greenwald, *et. al.*, (*J. Med. Chem.* 42:3657-3667 (1999)); and acetals and ketals of alcohol containing compounds (e.g., those obtained by condensation with chloromethyl methyl ether or chloromethyl ethyl ether according to methods known in the art).


The compounds of this invention may be prepared using methods known to those skilled in the art, or the novel methods of this invention. Specifically, the compounds of this invention with Formulae I-VII may be prepared as illustrated by the exemplary reaction in Scheme 1. Reaction of an acyl chloride with an amine in the presence a base such as triethylamine or *N,N*-diisopropylethylamine produces the amide.

Scheme 1



Alternatively, reaction of an acid with an amine in the presence of a coupling reagent such as EDC, or cyanuric chloride, also produces the amide as shown by the exemplary reaction in Scheme 2.

Scheme 2



An important aspect of the present invention is the discovery that compounds having Formula I-VII are activators of caspases and inducers of apoptosis. Therefore,

these compounds are useful in a variety of clinical conditions in which there is uncontrolled cell growth and spread of abnormal cells, such as in the case of cancer.

Another important aspect of the present invention is the discovery that compounds having Formula I-VII are potent and highly efficacious activators of caspases and inducers of apoptosis in drug resistant cancer cells, such as breast and prostate cancer cells, which enables these compounds to kill these drug resistant cancer cells. In comparison, most standard anti-cancer drugs are not effective in killing drug resistant cancer cells under the same conditions. Therefore, compounds of this invention are useful for the treatment of drug resistant cancer in animals.

The present invention includes a therapeutic method useful to modulate *in vivo* apoptosis or *in vivo* neoplastic disease, comprising administering to a subject in need of such treatment an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of the compound of Formulae I-VII, which functions as a caspase cascade activator and inducer of apoptosis.

The present invention also includes a therapeutic method comprising administering to an animal an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of said compound of Formulae I-VII, wherein said therapeutic method is useful to treat cancer, which is a group of diseases characterized by the uncontrolled growth and spread of abnormal cells. Such diseases include, but are not limited to, Hodgkin's disease, non-Hodgkin's lymphomas, acute lymphocytic leukemia, chronic lymphocytic leukemia, multiple myeloma, neuroblastoma, breast carcinomas, ovarian carcinomas, lung carcinomas, Wilms' tumor, cervical carcinomas, testicular carcinomas, soft-tissue sarcomas, primary macroglobulinemia, bladder carcinomas, chronic granulocytic leukemia, primary brain carcinomas, malignant melanoma, small-cell lung carcinomas, stomach carcinomas, colon carcinomas, malignant pancreatic insulinoma, malignant carcinoid carcinomas, malignant melanomas, choriocarcinomas, mycosis fungoides, head or neck carcinomas, osteogenic sarcoma, pancreatic carcinomas, acute granulocytic leukemia, hairy cell leukemia, neuroblastoma, rhabdomyosarcoma, Kaposi's sarcoma, genitourinary carcinomas, thyroid carcinomas, esophageal carcinomas, malignant hypercalcemia, cervical hyperplasia, renal cell carcinomas, endometrial carcinomas,

polycythemia vera, essential thrombocytosis, adrenal cortex carcinomas, skin cancer, and prostatic carcinomas.

In practicing the therapeutic methods, effective amounts of compositions containing therapeutically effective concentrations of the compounds formulated for oral, intravenous, local and topical application, for the treatment of neoplastic diseases and other diseases in which caspase cascade mediated physiological responses are implicated, are administered to an individual exhibiting the symptoms of one or more of these disorders. The amounts are effective to ameliorate or eliminate one or more symptoms of the disorders. An effective amount of a compound for treating a particular disease is an amount that is sufficient to ameliorate, or in some manner reduce, the symptoms associated with the disease. Such amount may be administered as a single dosage or may be administered according to a regimen, whereby it is effective. The amount may cure the disease but, typically, is administered in order to ameliorate the symptoms of the disease. Typically, repeated administration is required to achieve the desired amelioration of symptoms

In another embodiment, a pharmaceutical composition comprising a compound, or a pharmaceutically acceptable salt of said compound of Formulae I-VII, which functions as a caspase cascade activator and inducer of apoptosis in combination with a pharmaceutically acceptable vehicle is provided.

Another embodiment of the present invention is directed to a composition effective to inhibit neoplasia comprising a compound, or a pharmaceutically acceptable salt or prodrug of said compound of Formulae I-VII, which functions as a caspase cascade activator and inducer of apoptosis, in combination with at least one known cancer chemotherapeutic agent, or a pharmaceutically acceptable salt of said agent. Examples of known cancer chemotherapeutic agents which may be used for combination therapy include, but not are limited to alkylating agents such as busulfan, cis-platin, mitomycin C, and carboplatin; antimitotic agents such as colchicine, vinblastine, paclitaxel, and docetaxel; topo I inhibitors such as camptothecin and topotecan; topo II inhibitors such as doxorubicin and etoposide; RNA/DNA antimetabolites such as 5-azacytidine, 5-fluorouracil and methotrexate; DNA antimetabolites such as 5-fluoro-2'-deoxy-uridine, ara-C, hydroxyurea and thioguanine; antibodies such as Herceptin® and Rituxan®. Other

known cancer chemotherapeutic agents which may be used for combination therapy include melphalan, chlorambucil, cyclophosphamide, ifosfamide, vincristine, mitoguazone, epirubicin, acliarubicin, bleomycin, mitoxantrone, elliptinium, fludarabine, octreotide, retinoic acid, tamoxifen and alanosine.

5 In practicing the methods of the present invention, the compound of the invention may be administered together with at least one known chemotherapeutic agent as part of a unitary pharmaceutical composition. Alternatively, the compound of the invention may be administered apart from at least one known cancer chemotherapeutic agent. In one
10 embodiment, the compound of the invention and at least one known cancer chemotherapeutic agent are administered substantially simultaneously, i.e. the compounds are administered at the same time or one after the other, so long as the compounds reach therapeutic levels in the blood at the same time. On another embodiment, the compound of the invention and at least one known cancer chemotherapeutic agent are administered according to their individual dose schedule, so long as the compounds reach therapeutic
15 levels in the blood.

Another embodiment of the present invention is directed to a composition effective to inhibit neoplasia comprising a bioconjugates of said compound of Formulae I-VII, which functions as a caspase cascade activator and inducer of apoptosis, in bioconjugation with at least one known therapeutically useful antibody, such as Herceptin® or Rituxan®,
20 growth factors such as DGF, NGF, cytokines such as IL-2, IL-4, or any molecule that binds to the cell surface. The antibodies and other molecules will deliver the compound of Formulae I-VII to its targets and make it an effective anticancer agent. The bioconjugates could also enhance the anticancer effect of therapeutically useful antibodies, such as Herceptin® or Rituxan®.

25 Similarly, another embodiment of the present invention is directed to a composition effective in inhibiting neoplasia comprising a compound, or a pharmaceutically acceptable salt or prodrug of said compound of Formulae I-VII, which functions as a caspase cascade activator and inducer of apoptosis, in combination with radiation therapy. In this embodiment, the compound of the invention may be
30 administered at the same time as the radiation therapy is administered or at a different time.

Yet another embodiment of the present invention is directed to a composition effective for post-surgical treatment of cancer, comprising a compound, or a pharmaceutically acceptable salt or prodrug of said compound of Formulae I-VII, which functions as a caspase cascade activator and inducer of apoptosis. The invention also relates to a method of treating cancer by surgically removing the cancer and then treating the animal with one of the pharmaceutical compositions described herein.

A wide range of immune mechanisms operate rapidly following exposure to an infectious agent. Depending on the type of infection, rapid clonal expansion of the T and B lymphocytes occurs to combat the infection. The elimination of the effector cells following an infection is one of the major mechanisms for maintaining immune homeostasis. The elimination of the effector cells has been shown to be regulated by apoptosis. Autoimmune diseases have lately been determined to occur as a consequence of deregulated cell death. In certain autoimmune diseases, the immune system directs its powerful cytotoxic effector mechanisms against specialized cells such as oligodendrocytes in multiple sclerosis, the beta cells of the pancreas in diabetes mellitus, and thyrocytes in Hashimoto's thyroiditis (Ohsako, S. & Elkon, K.B., *Cell Death Differ.* 6:13-21 (1999)). Mutations of the gene encoding the lymphocyte apoptosis receptor Fas/APO-1/CD95 are reported to be associated with defective lymphocyte apoptosis and autoimmune lymphoproliferative syndrome (ALPS), which is characterized by chronic, histologically benign splenomegaly, generalized lymphadenopathy, hypergammaglobulinemia, and autoantibody formation. (Infante, A.J., *et al.*, *J. Pediatr.* 133:629-633 (1998) and Vaishnaw, A.K., *et al.*, *J. Clin. Invest.* 103:355-363 (1999)). It was reported that overexpression of Bcl-2, which is a member of the bcl-2 gene family of programmed cell death regulators with anti-apoptotic activity, in developing B cells of transgenic mice, in the presence of T cell dependent costimulatory signals, results in the generation of a modified B cell repertoire and in the production of pathogenic autoantibodies (Lopez-Hoyos, M., *et al.*, *Int. J. Mol. Med.* 1:475-483 (1998)). It is therefore evident that many types of autoimmune disease are caused by defects of the apoptotic process. One treatment strategy for such diseases is to turn on apoptosis in the lymphocytes that are causing the autoimmune disease (O'Reilly, L.A. & Strasser, A., *Inflamm. Res.* 48:5-21 (1999)).

Fas-Fas ligand (FasL) interaction is known to be required for the maintenance of immune homeostasis. Experimental autoimmune thyroiditis (EAT), characterized by autoreactive T and B cell responses and a marked lymphocytic infiltration of the thyroid, is a good model to study the therapeutic effects of FasL. Batteux, F., *et al.*, (*J. Immunol.* 162:603-608 (1999)) reported that by direct injection of DNA expression vectors encoding FasL into the inflamed thyroid, the development of lymphocytic infiltration of the thyroid was inhibited and induction of infiltrating T cells death was observed. These results show that FasL expression on thyrocytes may have a curative effect on ongoing EAT by inducing death of pathogenic autoreactive infiltrating T lymphocytes.

Bisindolylmaleimide VIII is known to potentiate Fas-mediated apoptosis in human astrocytoma 1321N1 cells and in Molt-4T cells; both of which were resistant to apoptosis induced by anti-Fas antibody in the absence of bisindolylmaleimide VIII. Potentiation of Fas-mediated apoptosis by bisindolylmaleimide VIII was reported to be selective for activated, rather than non-activated, T cells, and was Fas-dependent. Zhou T., *et al.*, (*Nat. Med.* 5:42-48 (1999)) reported that administration of bisindolylmaleimide VIII to rats during autoantigen stimulation prevented the development of symptoms of T cell-mediated autoimmune diseases in two models, the Lewis rat model of experimental allergic encephalitis and the Lewis adjuvant arthritis model. Therefore, the application of a Fas-dependent apoptosis enhancer such as bisindolylmaleimide VIII may be therapeutically useful for the more effective elimination of detrimental cells and inhibition of T cell-mediated autoimmune diseases. Therefore an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of the compound of Formulae I-VII, which functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for autoimmune diseases.

Psoriasis is a chronic skin disease that is characterized by scaly red patches. Psoralen plus ultraviolet A (PUVA) is a widely used and effective treatment for psoriasis vulgaris and Coven, *et al.*, *Photodermatol. Photoimmunol. Photomed.* 15:22-27 (1999), reported that lymphocytes treated with psoralen 8-MOP or TMP and UVA, displayed DNA degradation patterns typical of apoptotic cell death. Ozawa, *et al.*, *J. Exp. Med.* 189:711-718 (1999) reported that induction of T cell apoptosis could be the main mechanism by which 312-nm UVB resolves psoriasis skin lesions. Low doses of

methotrexate may be used to treat psoriasis to restore a clinically normal skin. Heenen, *et al.*, *Arch. Dermatol. Res.* 290:240-245 (1998), reported that low doses of methotrexate may induce apoptosis and that this mode of action could explain the reduction in epidermal hyperplasia during treatment of psoriasis with methotrexate. Therefore, an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of the compound of Formulae I-VII, which functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for psoriasis.

Synovial cell hyperplasia is a characteristic of patients with rheumatoid arthritis (RA). It is believed that excessive proliferation of RA synovial cells, as well as defects in synovial cell death, may be responsible for synovial cell hyperplasia. Wakisaka, *et al.*, *Clin. Exp. Immunol.* 114:119-128 (1998), found that although RA synovial cells could die via apoptosis through a Fas/FasL pathway, apoptosis of synovial cells was inhibited by proinflammatory cytokines present within the synovium. Wakisaka, *et al.*, also suggested that inhibition of apoptosis by the proinflammatory cytokines may contribute to the outgrowth of synovial cells, and lead to pannus formation and the destruction of joints in patients with RA. Therefore, an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of the compound of Formulae I-VII, which functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for rheumatoid arthritis.

There has been an accumulation of convincing evidence that apoptosis plays a major role in promoting resolution of the acute inflammatory response. Neutrophils are constitutively programmed to undergo apoptosis, thus limiting their pro-inflammatory potential and leading to rapid, specific, and non-phlogistic recognition by macrophages and semi-professional phagocytes (Savill, J., *J. Leukoc. Biol.* 61:375-380 (1997)). Boirivant, *et al.*, *Gastroenterology* 116:557-565 (1999), reported that lamina propria T cells, isolated from areas of inflammation in Crohn's disease, ulcerative colitis, and other inflammatory states, manifest decreased CD2 pathway-induced apoptosis. In addition, studies of cells from inflamed Crohn's disease tissue indicate that this defect is accompanied by elevated Bcl-2 levels. Therefore, an effective amount of a compound, or a pharmaceutically acceptable salt or prodrug of the compound of Formulae I-VII, which

functions as a caspase cascade activator and inducer of apoptosis, should be an effective treatment for inflammation.

Pharmaceutical compositions within the scope of this invention include all compositions wherein the compounds of the present invention are contained in an amount that is effective to achieve its intended purpose. While individual needs vary, determination of optimal ranges of effective amounts of each component is within the skill of the art. Typically, the compounds may be administered to mammals, e.g., humans, orally at a dose of 0.0025 to 50 mg/kg of body weight, per day, or an equivalent amount of the pharmaceutically acceptable salt thereof, to a mammal being treated for apoptosis-mediated disorders. Preferably, about 0.01 to about 10 mg/kg of body weight is orally administered to treat or prevent such disorders. For intramuscular injection, the dose is generally about one-half of the oral dose. For example, a suitable intramuscular dose would be about 0.0025 to about 25 mg/kg of body weight, and most preferably, from about 0.01 to about 5 mg/kg of body weight. If a known cancer chemotherapeutic agent is also administered, it is administered in an amount that is effective to achieve its intended purpose. The amounts of such known cancer chemotherapeutic agents effective for cancer are well known to those of skill in the art.

The unit oral dose may comprise from about 0.01 to about 50 mg, preferably about 0.1 to about 10 mg of the compound of the invention. The unit dose may be administered one or more times daily as one or more tablets each containing from about 0.1 to about 10, conveniently about 0.25 to 50 mg of the compound or its solvates.

In a topical formulation, the compound may be present at a concentration of about 0.01 to 100 mg per gram of carrier.

In addition to administering the compound as a raw chemical, the compounds of the invention may be administered as part of a pharmaceutical preparation containing suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the compounds into preparations which may be used pharmaceutically. Preferably, the preparations, particularly those preparations which may be administered orally and which may be used for the preferred type of administration, such as tablets, dragees, and capsules, and also preparations which may be administered rectally, such as suppositories, as well as suitable solutions for administration by injection

or orally, contain from about 0.01 to 99 percent, preferably from about 0.25 to 75 percent of active compound(s), together with the excipient.

Also included within the scope of the present invention are the non-toxic pharmaceutically acceptable salts of the compounds of the present invention. Acid addition salts are formed by mixing a solution of the particular apoptosis inducers of the present invention with a solution of a pharmaceutically acceptable non-toxic acid such as hydrochloric acid, fumaric acid, maleic acid, succinic acid, acetic acid, citric acid, tartaric acid, carbonic acid, phosphoric acid, oxalic acid, and the like. Basic salts are formed by mixing a solution of the particular apoptosis inducers of the present invention with a solution of a pharmaceutically acceptable non-toxic base such as sodium hydroxide, potassium hydroxide, choline hydroxide, sodium carbonate, Tris, N-methyl-glucamine and the like.

The pharmaceutical compositions of the invention may be administered to any animal which may experience the beneficial effects of the compounds of the invention. Foremost among such animals are mammals, e.g., humans and veterinary animals, although the invention is not intended to be so limited.

The pharmaceutical compositions of the present invention may be administered by any means that achieve their intended purpose. For example, administration may be by parenteral, subcutaneous, intravenous, intramuscular, intraperitoneal, transdermal, buccal, intrathecal, intracranial, intranasal or topical routes. Alternatively, or concurrently, administration may be by the oral route. The dosage administered will be dependent upon the age, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired.

The pharmaceutical preparations of the present invention are manufactured in a manner which is itself known, for example, by means of conventional mixing, granulating, dragee-making, dissolving, or lyophilizing processes. Thus, pharmaceutical preparations for oral use may be obtained by combining the active compounds with solid excipients, optionally grinding the resulting mixture and processing the mixture of granules, after adding suitable auxiliaries, if desired or necessary, to obtain tablets or dragee cores.

Suitable excipients are, in particular, fillers such as saccharides, for example lactose or sucrose, mannitol or sorbitol, cellulose preparations and/or calcium phosphates,

for example tricalcium phosphate or calcium hydrogen phosphate, as well as binders such as starch paste, using, for example, maize starch, wheat starch, rice starch, potato starch, gelatin, tragacanth, methyl cellulose, hydroxypropylmethylcellulose, sodium carboxymethylcellulose, and/or polyvinyl pyrrolidone. If desired, disintegrating agents may be added such as the above-mentioned starches and also carboxymethyl-starch, cross-linked polyvinyl pyrrolidone, agar, or alginic acid or a salt thereof, such as sodium alginate. Auxiliaries are, above all, flow-regulating agents and lubricants, for example, silica, talc, stearic acid or salts thereof, such as magnesium stearate or calcium stearate, and/or polyethylene glycol. Dragee cores are provided with suitable coatings which, if desired, are resistant to gastric juices. For this purpose, concentrated saccharide solutions may be used, which may optionally contain gum arabic, talc, polyvinyl pyrrolidone, polyethylene glycol and/or titanium dioxide, lacquer solutions and suitable organic solvents or solvent mixtures. In order to produce coatings resistant to gastric juices, solutions of suitable cellulose preparations such as acetylcellulose phthalate or hydroxypropymethylcellulose phthalate, are used. Dye stuffs or pigments may be added to the tablets or dragee coatings, for example, for identification or in order to characterize combinations of active compound doses.

Other pharmaceutical preparations which may be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a plasticizer such as glycerol or sorbitol. The push-fit capsules may contain the active compounds in the form of granules which may be mixed with fillers such as lactose, binders such as starches, and/or lubricants such as talc or magnesium stearate and, optionally, stabilizers. In soft capsules, the active compounds are preferably dissolved or suspended in suitable liquids, such as fatty oils, or liquid paraffin. In addition, stabilizers may be added.

Possible pharmaceutical preparations which may be used rectally include, for example, suppositories, which consist of a combination of one or more of the active compounds with a suppository base. Suitable suppository bases are, for example, natural or synthetic triglycerides, or paraffin hydrocarbons. In addition, it is also possible to use gelatin rectal capsules which consist of a combination of the active compounds with a base. Possible base materials include, for example, liquid triglycerides, polyethylene glycols, or paraffin hydrocarbons.

Suitable formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form, for example, water-soluble salts and alkaline solutions. In addition, suspensions of the active compounds as appropriate oily injection suspensions may be administered. Suitable lipophilic solvents or vehicles include fatty oils, for example, sesame oil, or synthetic fatty acid esters, for example, ethyl oleate or triglycerides or polyethylene glycol-400 (the compounds are soluble in PEG-400) or cremophor, or cyclodextrins. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension include, for example, sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension may also contain stabilizers.

In accordance with one aspect of the present invention, compounds of the invention are employed in topical and parenteral formulations and are used for the treatment of skin cancer.

The topical compositions of this invention are formulated preferably as oils, creams, lotions, ointments, gels and the like by choice of appropriate carriers. Suitable carriers include vegetable or mineral oils, white petrolatum (white soft paraffin), branched chain fats or oils, animal fats and high molecular weight alcohol (greater than C_{12}). The preferred carriers are those in which the active ingredient is soluble. Emulsifiers, stabilizers, humectants and antioxidants may also be included as well as agents imparting color or fragrance, if desired. Additionally, transdermal penetration enhancers may be employed in these topical formulations. Examples of such enhancers are found in U.S. Patent Nos. 3,989,816 and 4,444,762.

Creams are preferably formulated from a mixture of mineral oil, self-emulsifying beeswax and water in which mixture the active ingredient, dissolved in a small amount of an oil such as almond oil, is admixed. A typical example of such a cream is one which includes about 40 parts water, about 20 parts beeswax, about 40 parts mineral oil and about 1 part almond oil.

Ointments may be formulated by mixing a solution of the active ingredient in a vegetable oil such as almond oil with warm soft paraffin and allowing the mixture to cool. A typical example of such an ointment is one which includes about 30% almond oil and about 70% white soft paraffin by weight.

The following examples are illustrative, but not limiting, of the method and compositions of the present invention. Other suitable modifications and adaptations of the variety of conditions and parameters normally encountered in clinical therapy and which are obvious to those skilled in the art are within the spirit and scope of the invention.

5

Example 1

6-Chloro-N-(2-nitro-4-trifluoromethylphenyl)-3-pyridinecarboxamide

To a mixture of 6-chloronicotinoyl chloride (100 mg, 0.6 mmol) and 2-nitro-4-trifluoromethylaniline (117 mg, 0.6 mmol) in THF (10 ml) was added triethylamine (0.16 ml, 1.1 mmol). The mixture was stirred at room temperature overnight and then diluted with ethyl acetate. Water was added to dissolve the precipitate and the organic phase was separated and washed with water, then dried over anhydrous sodium sulfate. The product was purified by column chromatography to give the title compound as a pale yellow solid.

¹H NMR (DMSO-d₆): 11.27 (s, 1H), 8.97 (d, J = 2.7, 1H), 8.38-8.35 (m, 2H), 8.19 (d, J = 8.4, 1H), 7.96 (d, J = 8.7, 1H), 7.78 (d, J = 8.1, 1H).

15

Example 2

6-Chloro-N-(4-chloro-2-nitrophenyl)-3-pyridinecarboxamide

20

The title compound was prepared from 6-chloronicotinoyl chloride and 4-chloro-2-nitroaniline and was obtained as a yellow solid as described in Example 1. ¹H NMR (DMSO-d₆): 11.31 (s, 1H), 9.02 (d, J = 2.4, 1H), 8.96 (d, J = 9.3, 1H), 8.31 (d, J = 2.4, 1H), 8.24-8.20 (m, 1H), 7.73-7.69 (m, 1H), 7.54 (d, J = 8.4, 1H).

25

Example 3

6-Chloro-N-(4-fluoro-2-nitrophenyl)-3-pyridinecarboxamide

The title compound was prepared from 6-chloronicotinoyl chloride and 4-fluoro-2-nitroaniline and was obtained as a yellow solid as described in Example 1. ¹H NMR

30

-34-

(DMSO- d_6): 10.97 (s, 1H), 8.94 (d, $J = 2.7$, 1H), 8.36-8.32 (m, 1H), 8.02-7.98 (m, 1H), 7.78-7.75 (m, 1H), 7.73-7.72 (m, 1H), 7.70 (d, $J = 1.5$, 1H).

Example 4

5 *6-Chloro-N-(3-bromo-4-methoxy-6-nitrophenyl)-3-pyridinecarboxamide*

The title compound was prepared from 6-chloronicotinoyl chloride and 3-bromo-4-methoxy-6-nitroaniline and was obtained as a yellow solid as described in Example 1. ^1H NMR (DMSO- d_6): 10.84 (s, 1H), 8.93 (d, $J = 2.4$, 1H), 8.34-8.30 (m, 1H), 7.98 (s, 1H),
10 7.76 (d, $J = 8.4$, 1H), 7.70 (s, 1H), 3.97 (s, 3H).

Example 5

6-Chloro-N-(2,4-dimethoxyphenyl)-3-pyridinecarboxamide

15 The title compound was prepared from 6-chloronicotinoyl chloride and 2,4-dimethoxyaniline as described in Example 1. The reaction mixture was filtered to remove the precipitate and the filtrate was evaporated. The resulting solid was purified by CH_2Cl_2 /hexane recrystallization to give a light purple solid in a yield of 59%. ^1H NMR (DMSO- d_6): 9.77 (s, 1H), 8.93 (s, 1H), 8.35-8.31 (m, 1H), 7.68 (d, $J = 7.8$, 1H), 7.46 (d, $J = 9.0$, 1H), 6.67 (d, $J = 2.4$, 1H), 6.57-6.53 (m, 1H), 3.80 (s, 3H), 3.78 (s, 3H).
20

Example 6

2-Chloro-N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide

25 The title compound was prepared from 2-chloronicotinoyl chloride and 4-methoxy-2-nitroaniline and was obtained as a yellow solid as described in Example 1. ^1H NMR (CDCl_3): 10.76 (s, 1H), 8.78 (d, $J = 9.3$, 1H), 8.58-8.55 (m, 1H), 8.13-8.09 (m, 1H), 7.73 (d, $J = 3.0$, 1H), 7.45-7.40 (m, 1H), 7.34-7.28 (m, 1H), 3.90 (s, 3H).

Example 7***6-Chloro-N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinoyl chloride and 4-methoxy-2-nitroaniline and was obtained as a yellow solid as described in Example 1. ¹H NMR (CDCl₃): 11.13 (s, 1H), 9.01 (d, J = 2.7, 1H), 8.83 (d, J = 9.3, 1H), 8.24-8.20 (m, 1H), 7.76 (d, J = 2.7, 1H), 7.52 (d, J = 8.7, 1H), 7.35-7.31 (m, 1H), 3.90 (s, 3H).

Example 8***6-Chloro-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinoyl chloride and 4-ethoxy-2-nitroaniline and was obtained as a yellow solid as described in Example 1. ¹H NMR (CDCl₃): 11.13 (s, 1H), 9.01 (d, J = 2.4, 1H), 8.82 (d, J = 9.3, 1H), 8.24-8.20 (m, 1H), 7.74 (d, J = 3.0, 1H), 7.51 (d, J = 7.8, 1H), 7.34-7.30 (m, 1H), 4.11 (q, J = 6.9, 2H), 1.47 (t, J = 6.9, 3H).

Example 9***6-Chloro-N-(4-methyl-2-nitrophenyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinoyl chloride and 4-methyl-2-nitroaniline and was obtained as a yellow solid as described in Example 1. ¹H NMR (CDCl₃): 11.29 (s, 1H), 9.02 (d, J = 2.7, 1H), 8.81 (d, J = 8.7, 1H), 8.24-8.21 (m, 1H), 8.11 (s, 1H), 7.58-7.50 (m, 2H), 2.44 (s, 3H).

Example 10***6-Chloro-N-(2-nitrophenyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinoyl chloride and 2-nitroaniline and was obtained as a yellow solid as described in Example 1. ¹H NMR

(CDCl₃): 10.99 (s, 1H), 8.95-8.94 (m, 1H), 8.36-8.33 (m, 1H), 8.02-7.99 (m, 1H), 7.80-7.69 (m, 3H), 7.48-7.42 (m, 1H).

Example 11

6-Chloro-N-(4-methoxyphenyl)-3-pyridinecarboxamide

The title compound was prepared from 6-chloronicotinoyl chloride and *p*-anisidine and was obtained as a white solid as described in Example 1. ¹H NMR (CDCl₃): 8.85 (d, J = 2.1, 1H), 8.19-8.15 (m, 1H), 7.65 (s, 1H), 7.54-7.46 (m, 3H), 6.96-6.91 (m, 2H), 3.83 (s, 3H).

Example 12

2,6-Dichloro-N-(4-methoxy-2-nitrophenyl)-4-pyridinecarboxamide

The title compound was prepared from 2,6-dichloropyridine-4-carbonyl chloride and 4-methoxy-2-nitroaniline and was obtained as a yellow solid as described in Example 1. ¹H NMR (CDCl₃): 11.12 (s, 1H), 8.78 (d, J = 9.3, 1H), 7.77 (d, J = 3.0, 1H), 7.75 (s, 2H), 7.35-7.31 (m, 1H), 3.91 (s, 3H).

Example 13

2-Chloro-4-trifluoromethyl-N-(4-methoxy-2-nitrophenyl)-5-pyrimidinecarboxamide

The title compound was prepared from 2-chloro-4-(trifluoromethyl)pyrimidine-5-carbonyl chloride and 4-methoxy-2-nitroaniline and was obtained as a light yellow solid as described in Example 1. ¹H NMR (CDCl₃): 10.52 (s, 1H), 9.04 (s, 1H), 8.68 (d, J = 9.3, 1H), 7.76 (d, J = 3.0, 1H), 7.35-7.31 (m, 1H), 3.91 (s, 3H).

Example 14***N-(4-methoxy-2-nitrophenyl)-4-pyridinecarboxamide***

The title compound was prepared from isonicotinoyl chloride and 4-methoxy-2-nitroaniline and was obtained as an orange yellow solid as described in Example 1. ¹H NMR (CDCl₃): 10.79 (s, 1H), 8.83-8.81 (m, 2H), 7.85-7.83 (m, 2H), 7.61-7.55 (m, 2H), 7.40-7.36 (m, 1H), 3.87 (s, 3H).

Example 15***2,6-Dichloro-N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide***

To a solution of 2,6-dichloronicotinic acid (100 mg, 0.52 mmol) in CH₂Cl₂ (10 mL) was added cyanuric chloride (96 mg, 0.52 mmol). The mixture was stirred for half an hour, then 4-methoxy-2-nitroaniline (88 mg, 0.52 mmol) and triethylamine (0.1 mL, 0.72 mmol) were added. The resulting mixture was stirred at room temperature overnight and then diluted with ethyl acetate, washed with water. The organic phase was separated, dried over anhydrous Na₂SO₄, concentrated and purified by chromatography to give the title compound (40 mg, 22%) as a yellow solid. ¹H NMR (DMSO-*d*₆): 10.84 (bs, 1H), 8.09 (d, J = 7.8, 1H), 7.75 (d, J = 7.8, 1H), 7.57-7.52 (m, 2H), 7.37-7.33 (m, 1H), 3.86 (s, 3H).

Example 16***5,6-Dichloro-N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide***

The title compound was prepared from 5,6-dichloronicotinic acid and 4-methoxy-2-nitroaniline and was obtained as a yellow solid as described in Example 15. ¹H NMR (DMSO-*d*₆): 10.85 (bs, 1H), 8.88 (d, J = 2.4, 1H), 8.59 (d, J = 2.4, 1H), 7.58-7.54 (m, 2H), 7.39-7.35 (m, 1H), 3.87 (s, 3H).

Example 17***6-Chloro-N-(2,4-dinitrophenyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinoyl chloride and 2,4-dinitroaniline and was obtained as a yellow solid as described in Example 1. ¹H NMR (CDCl₃): 11.69 (bs, 1H), 9.28-9.23 (m, 2H), 9.06-9.05 (m, 1H), 8.61-8.57 (m, 1H), 8.28-8.24 (m, 1H), 7.59-7.56 (m, 1H).

Example 18***4-Chloro-N-(4-ethoxy-2-nitrophenyl)-benzoylamide***

The title compound was prepared from 4-chlorobenzoyl chloride and 4-ethoxy-2-nitroaniline and was obtained as a yellow solid as described in Example 1. ¹H NMR (CDCl₃): 11.10 (bs, 1H), 8.86 (d, J = 9.3, 1H), 7.93 (d, J = 8.4, 2H), 7.73 (d, J = 3.0, 1H), 7.51 (d, J = 8.4, 2H), 7.32-7.27 (m, 1H), 4.11 (q, J = 6.9, 2H), 1.46 (t, J = 6.9, 3H).

Example 19***6-Chloro-N-(4,5-difluoro-2-nitrophenyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinoyl chloride and 4,5-difluoro-2-nitroaniline and was obtained as a yellow solid as described in Example 1. ¹H NMR (CDCl₃): 11.49 (bs, 1H), 9.04-8.97 (m, 2H), 8.25-8.19 (m, 2H), 7.56-7.53 (m, 1H).

Example 20***6-Chloro-N-(4-methoxy-2-methylphenyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinoyl chloride and 4-methoxy-2-methylaniline and was obtained as a white solid as described in Example 1. ¹H NMR (CDCl₃): 8.88 (bs, 1H), 8.19 (d, J = 7.2, 1H), 7.61-7.47 (m, 3H), 6.81-6.79 (m, 2H), 3.81 (s, 3H), 2.30 (s, 3H).

Example 21***6-Chloro-N-(4-cyano-2-nitrophenyl)-3-pyridinecarboxamide***

5 The title compound was prepared from 6-chloronicotinoyl chloride and 4-amino-3-nitrobenzonitrile and was obtained as a light yellow solid as described in Example 1. ¹H NMR (CDCl₃): 11.58 (s, 1H), 9.18 (d, J = 8.7, 1H), 9.04 (d, J = 2.4, 1H), 8.65 (d, J = 2.1, 1H), 8.26-8.22 (m, 1H), 8.00-7.96 (m, 1H), 7.56 (d, J = 9.0, 1H).

Example 22***6-Fluoro-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide***

10 To the solution of KMnO₄ (284 mg, 1.8 mmol) in H₂O (50 mL) was added 2-fluoro-5-methylpyridine (100 mg, 0.9 mmol) at room temperature. The reaction mixture was then heated to 100°C for 4 h. The resulting precipitate was filtered and discarded. 15 The aqueous solution was washed with 3:2, hexane: ethyl acetate (2 x 20 mL), acidified with 2N HCl, then extracted with ethyl acetate (3 x 20 mL). The organic extracts were combined and washed with H₂O and brine, dried over anhydrous Na₂SO₄. The solvent was evaporated *in vacuo* to give 6-fluoronicotinic acid (30 mg) as a white solid. 20 The title compound was prepared from 6-fluoronicotinic acid and 4-ethoxy-2-nitroaniline as a yellow solid as described in Example 15. ¹H NMR (CDCl₃): 11.09 (bs, 1H), 8.88 (d, J = 2.4, 1H), 8.81 (d, J = 9.3, 1H), 8.42-8.35 (m, 1H), 7.74 (d, J = 3.3, 1H), 7.33-7.29 (m, 1H), 7.13-7.10 (m, 1H), 4.11 (q, J = 7.2, 2H), 1.47 (t, J = 7.2, 3H).

Example 23***6-Chloro-N-(2-nitro-4-trifluoromethoxyphenyl)-3-pyridinecarboxamide***

25 The title compound was prepared from 6-chloronicotinoyl chloride and 2-nitro-4-(trifluoromethoxy)aniline and was obtained as a yellow solid as described in Example 1.

¹H NMR (CDCl₃): 11.33 (bs, 1H), 9.06 (d, J = 9.3, 1H), 9.02 (d, J = 2.7, 1H), 8.25-8.21 (m, 1H), 8.19 (d, J = 2.7, 1H), 7.65-7.61 (m, 1H), 7.54 (d, J = 8.4, 1H).

Example 24

6-Chloro-N-(4-benzoyl-2-nitrophenyl)-3-pyridinecarboxamide

The title compound was prepared from 6-chloronicotinoyl chloride and 4-amino-3-nitrobenzophenone and was obtained as a yellow solid as described in Example 1. ¹H NMR (CDCl₃): 11.61 (bs, 1H), 9.12 (d, J = 8.7, 1H), 9.06 (d, J = 2.4, 1H), 8.79 (d, J = 1.8, 1H), 8.28-8.25 (m, 1H), 8.23-8.19 (m, 1H), 7.82-7.81 (m, 1H), 7.80-7.79 (m, 1H), 7.67-7.64 (m, 1H), 7.58-7.53 (m, 3H).

Example 25

6-Chloro-N-(4-benzyloxy-2-nitrophenyl)-3-pyridinecarboxamide

The title compound was prepared from 6-chloronicotinoyl chloride and 4-benzyloxy-2-nitroaniline and was obtained as a yellow solid as described in Example 1. ¹H NMR (CDCl₃): 11.13 (bs, 1H), 9.00 (d, J = 3.0, 1H), 8.83 (d, J = 9.3, 1H), 8.23-8.20 (m, 1H), 7.85 (d, J = 3.0, 1H), 7.53-7.46 (m, 1H), 7.44-7.37 (m, 6H), 5.15 (s, 2H).

Example 26

6-Methyl-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide

The title compound was prepared from 6-methylnicotinic acid and 4-ethoxy-2-nitroaniline and was obtained as a yellow solid as described in Example 15. ¹H NMR (CDCl₃): 11.09 (s, 1H), 9.12 (d, J = 2.1, 1H), 8.85 (d, J = 9.3, 1H), 8.16-8.13 (m, 1H), 7.73 (d, J = 3.0, 1H), 7.34-7.29 (m, 2H), 4.11 (q, J = 6.9, 2H), 2.67 (s, 3H), 1.46 (t, J = 6.9, 3H).

Example 27***6-Chloro-N-(4,5-dimethyl-2-nitrophenyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinoyl chloride and 4,5-dimethyl-2-nitroaniline and was obtained as an orange solid as described in Example 1. ¹H NMR (CDCl₃): 11.38 (s, 1H), 9.03 (d, J = 2.1, 1H), 8.72 (s, 1H), 8.24-8.21 (m, 1H), 8.07 (s, 1H), 7.52 (d, J = 8.7, 1H), 2.41 (s, 3H), 2.34 (s, 3H).

Example 28***6-Chloro-N-(4-methoxy-2-nitrophenyl)-1-N-oxide-3-pyridinecarboxamide***

To a stirred solution of 30% H₂O₂ (2 mL) was added trifluoroacetic anhydride (1 mL) dropwise at 0°C. The resulting solution was stirred at 0°C for 30 min, then 6-chloro-N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide (30 mg, 0.0975 mmol) was added. The mixture was stirred at 90°C for 30 min. The resulting yellow solution was cooled in an ice bath for 1 h. The precipitate was filtered, washed with H₂O and MeOH, and dried to give 15 mg (0.0463 mmol, 47%) of the title compound as a yellow powder. ¹H NMR (CDCl₃): 11.11 (s, 1H), 8.95-8.76 (m, 1H), 8.78 (d, J = 9.3, 1H), 7.76 (d, J = 3.0, 1H), 7.66-7.65 (m, 2H), 7.35-7.31 (m, 1H), 3.91 (s, 3H).

Example 29***N-(4-Methoxy-2-nitrophenyl)-3-pyridinecarboxamide***

The title compound was prepared from nicotinoyl chloride and 4-methoxy-2-nitroaniline and was obtained as an orange solid as described in Example 1. ¹H NMR (DMSO-d₆): 10.74 (s, 1H), 9.10 (d, J = 2.4, 1H), 8.80-8.78 (m, 1H), 7.62-7.54 (m, 3H), 7.39-7.35 (m, 1H), 3.87 (s, 3H).

Example 30***6-Chloro-N-(4-ethoxy-2-nitrophenyl)-N-methyl-3-pyridinecarboxamide***

To a solution of 6-chloro-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide (50 mg, 0.155 mmol) in DMF (2 ml) was added sodium methoxide (12 mg, 0.222 mmol) in DMF (1 ml) and the reaction mixture was stirred for 10 min. To the mixture was added iodomethane (22 mg, 0.155 mmol) slowly and the mixture was stirred for at room temperature for 3 h. It was diluted with ethyl acetate and washed with water and brine, dried over anhydrous sodium sulfate, and evaporated. The residue was purified by column chromatography to yield the titled compound as a light brown liquid. ¹H NMR (CDCl₃): 8.17 (d, J = 2.1, 1H), 7.68-7.65 (m, 1H), 7.35 (d, J = 2.7, 1H), 7.21 (t, J = 7.8, 2H), 7.10-7.06 (m, 1H), 4.08-4.01 (m, 2H), 3.40 (s, 3H), 1.44 (t, J = 6.9, 3H).

Example 31***2,5-Dimethyl-N-(4-ethoxy-2-nitrophenyl)-3-furancarboxamide***

The title compound was prepared from 2,5-dimethyl-3-furoic acid and 4-ethoxy-2-nitroaniline as an orange solid as described in Example 15. ¹H NMR (CDCl₃): 10.59 (s, 1H), 8.83 (d, J = 9.0, 1H), 7.70 (d, J = 3.0, 1H), 7.27-7.25 (m, 1H), 6.24 (s, 1H), 4.08 (q, J = 6.9, 2H), 2.62 (s, 3H), 2.30 (s, 3H), 1.45 (t, J = 6.9, 3H).

Example 32***6-Chloro-N-(5-bromo-3-nitro-2-pyridyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinoyl chloride and 2-amino-5-bromo-3-nitropyridine as a light yellow solid as described in Example 1. ¹H NMR (CDCl₃): 10.56 (s, 1H), 9.00 (d, J = 2.7, 1H), 8.82 (d, J = 2.1, 1H), 8.69 (d, J = 2.1, 1H), 8.27-8.23 (m, 1H), 7.53 (d, J = 8.1, 1H).

Example 33***N-(4-Ethoxy-2-nitrophenyl)-2-pyrazinecarboxamide***

The title compound was prepared from 2-pyrazinecarboxylic acid and 4-ethoxy-2-nitroaniline as a yellow solid as described in Example 15. ¹H NMR (CDCl₃): 12.34 (s, 1H), 9.50 (d, J = 1.2, 1H), 8.91 (d, J = 9.0, 1H), 8.84 (d, J = 2.4, 1H), 8.72-8.70 (m, 1H), 7.75 (d, J = 3.0, 1H), 7.33-7.28 (m, 1H), 4.12 (q, J = 6.9, 2H), 1.47 (t, J = 6.9, 3H).

Example 34***5-Methyl-N-(4-ethoxy-2-nitrophenyl)-2-pyrazinecarboxamide***

The title compound was prepared from 2-methylpyrazine-5-carboxylic acid and 4-ethoxy-2-nitroaniline as a yellow solid as described in Example 15. ¹H NMR (CDCl₃): 12.82 (s, 1H), 9.36 (s, 1H), 8.90 (d, J = 9.6, 1H), 8.56 (d, J = 1.0, 1H), 7.74 (d, J = 2.7, 1H), 7.32-7.29 (m, 1H), 4.11 (q, J = 6.9, 2H), 2.71 (s, 3H), 1.46 (t, J = 6.9, 3H).

Example 35***6-Chloro-N-(4-chloro-2-cyanophenyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinoyl chloride and 2-amino-5-chlorobenzonitrile as a white solid as described in Example 1. ¹H NMR (CDCl₃): 8.96 (d, J = 2.7, 1H), 8.53-8.50 (M, 1H), 8.24 (s, 1H), 8.19-8.15 (m, 1H), 7.67-7.63 (m, 2H), 7.54-7.51 (m, 1H).

Example 36***6-Chloro-N-(4-chloro-2-amidophenyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinoyl chloride and 2-amino-5-chlorobenzamide as a white solid as described in Example 1. ¹H NMR (DMSO-*d*₆): 8.91

(d, J = 2.7, 1H), 8.62 (d, J = 9.3, 1H), 8.55 (s, 1H), 8.30-8.27 (m, 1H), 8.04 (s, 1H), 8.01 (d, J = 2.7, 1H), 7.78 (d, J = 8.4, 1H), 7.69-7.66 (m, 1H).

Example 37

6-Trifluoromethyl-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide

The title compound was prepared from 6-trifluoromethylnicotinic acid and 4-ethoxy-2-nitroaniline as a yellow solid as described in Example 15. ¹H NMR (CDCl₃): 11.23 (s, 1H), 9.32 (d, J = 3.6, 1H), 8.83 (d, J = 9.3, 1H), 8.47-8.43 (m, 1H), 7.88 (d, J = 8.1, 1H), 7.76 (d, J = 3.0, 1H), 7.35-7.31 (m, 1H), 4.12 (q, J = 6.9, 2H), 1.47 (t, J = 6.9, 3H).

Example 38

6-Chloro-N-(4-chloro-2-methoxycarbonylphenyl)-3-pyridinecarboxamide

The title compound was prepared from 6-chloronicotinoyl chloride and methyl 2-amino-5-chlorobenzoate as a white solid as described in Example 1. ¹H NMR (CDCl₃): 12.13 (s, 1H), 9.06 (d, J = 3.0, 1H), 8.86 (d, J = 9.3, 1H), 8.29-8.25 (m, 1H), 8.08 (d, J = 2.4, 1H), 7.60-7.57 (m, 1H), 7.50 (d, J = 8.4, 1H), 3.99 (s, 3H).

Example 39

5-Bromo-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide

The title compound was prepared from 5-bromonicotinic acid and 4-ethoxy-2-nitroaniline as a yellow solid as described in Example 15. ¹H NMR (CDCl₃): 11.11 (s, 1H), 9.11 (d, J = 2.1, 1H), 8.88 (d, J = 1.8, 1H), 8.80 (d, J = 9.3, 1H), 8.42 (t, J = 2.1, 1H), 7.75 (d, J = 2.7, 1H), 7.34-7.30 (m, 1H), 4.12 (q, J = 6.9, 2H), 1.47 (t, J = 6.9, 3H).

Example 40***N-(4-Ethoxy-2-nitrophenyl)-2-pyridinecarboxamide N-oxide***

The title compound was prepared from picolinic acid N-oxide and 4-ethoxy-2-nitroaniline as a yellow solid as described in Example 15. ¹H NMR (DMSO-*d*₆): 8.57-8.54 (m, 1H), 8.40 (d, J = 9.0, 1H), 8.37-8.33 (m, 1H), 7.76-7.75 (m, 2H), 7.61 (d, J = 3.0, 1H), 7.43-7.39 (m, 1H), 4.14 (q, J = 6.9, 2H), 1.36 (t, J = 6.9, 3H).

Example 41***4-Amino-3-nitrophenyl 6-Chloro-3-pyridinecarboxylate (A) and 6-Chloro-N-(4-hydroxy-2-nitrophenyl)-3-pyridinecarboxamide (B)***

A mixture of 4-amino-3-nitrophenol (200 mg, 1.30 mmol), 6-chloronicotinoyl chloride (229 mg, 1.30 mmol) and N,N-diisopropylethylamine (200 ul) in THF (5 ml) was stirred overnight at room temperature. The resulting solid was collected by filtration, wash with hexane:ethyl acetate (1:1) and dried under vacuo to give compound A as an orange solid (14 mg). ¹H NMR (DMSO-*d*₆): 9.08 (dd, J = 8.0, 2.6 Hz, 1H), 8.47 (dd, J = 2.6, 8.3 Hz, 1H), 7.97 (d, J = 2.7 Hz, 1H), 7.78 (dd, J = 0.8, 8.3 Hz, 1H), 7.55 (s, 2H), 7.45 (dd, J = 2.7, 9 Hz, 1H), 7.10 (d, J = 9.0 Hz, 1H). The filtrate was diluted with ethyl acetate (50 ml), washed with water (25 ml), aqueous saturated NaCl (25 ml) and dried over anhydrous sodium sulfate. The solution was concentrated in vacuo and the product was purified by column chromatography using hexane:ethyl acetate (3:1) to give the product B as a dark yellow solid (9 mg). ¹H NMR (DMSO-*d*₆): 9.13 (d, J = 2.4 Hz, 1H), 8.96 (d, J = 2.4 Hz, 1H), 8.52 (dd, J = 1.8, 8.7 Hz, 1H), 8.36 (dd, J = 2.4, 8.4 Hz, 1H), 8.07 (s, 1H), 7.85-7.80 (m, 2H), 7.73 (d, J = 8.4 Hz, 1H).

Example 42***N-(4-Ethoxy-2-nitrophenyl)-3-pyrrolecarboxamide***

The title compound was prepared from pyrrole-3-carboxylic acid and 4-ethoxy-2-nitroaniline as a yellow solid as described in Example 15. ¹H NMR (CDCl₃): 10.82 (s, 1H), 8.88 (d, J = 9.3, 1H), 8.65 (s, 1H), 7.70 (d, J = 3.0, 1H), 7.54-7.52 (m, 1H), 7.28-7.24 (m, 1H), 6.87-6.85 (m, 1H), 6.71-6.69 (m, 1H), 4.08 (q, J = 6.9, 2H), 1.45 (t, J = 6.9, 3H).

Example 43***6-Chloro-N-(2-nitro-5-imidazolyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinonyl chloride and 5-amino-4-nitroimidazole as a brown solid as described in Example 1. ¹H NMR (DMSO-*d*₆): 8.97-8.96 (m, 1H), 8.39-8.36 (m, 1H), 7.79-7.76 (m, 1H), 7.68 (s, 1H), 7.37 (s, 1H).

Example 44***6-Chloro-N-(4-t-butyl-2-nitrophenyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinic chloride and 4-(t-butyl)-2-nitroaniline as a yellow solid as described in Example 1. ¹H NMR (CDCl₃): 11.30 (s, 1H), 9.03 (d, J = 2.1, 1H), 8.83 (d, J = 9.0, 1H), 8.29 (d, J = 2.7, 1H), 8.25-8.21 (m, 1H), 7.80-7.76 (m, 1H), 7.52 (d, J = 8.4, 1H), 1.38 (s, 9H).

Example 45***N-(4-Ethoxy-2-nitrophenyl)-5-pyrimidinecarboxamide***

The title compound was prepared from pyrimidine-5-carboxylic acid and 4-ethoxy-2-nitroaniline as a yellow solid as described in Example 15. ¹H NMR (CDCl₃): 11.16 (s, 1H), 9.43 (s, 1H), 9.32 (s, 2H), 8.81 (d, J = 9.3, 1H), 7.76 (d, J = 3.0, 1H), 7.34-7.27 (m, 1H), 4.12 (q, J = 6.9, 2H), 1.47 (t, J = 6.9, 3H).

Example 46***N-(4-Ethoxy-2-nitrophenyl)-2-pyridinecarboxamide***

5 The title compound was prepared from picolinic acid and 4-ethoxy-2-nitroaniline as a yellow solid as described in Example 15. ¹H NMR (CDCl₃): 12.54 (s, 1H), 8.92 (d, J = 9.3, 1H), 8.73 (d, J = 3.9, 1H), 8.28 (d, J = 6.9, 1H), 7.95-7.90 (m, 1H), 7.74 (d, J = 3.0, 1H), 7.54-7.50 (m, 1H), 7.31-7.27 (m, 1H), 4.11 (q, J = 6.9, 2H), 1.46 (t, J = 6.9, 3H).

Example 47***6-Chloro-N-(2-cyano-4,5-dimethoxyphenyl)-3-pyridinecarboxamide***

10 The title compound was prepared from 6-chloronicotinoyl chloride and 2-amino-4,5-dimethoxybenzonitrile as a white solid as described in Example 1. ¹H NMR (CDCl₃): 8.97 (d, J = 2.7, 1H), 8.18-8.15 (m, 3H), 7.52 (d, J = 8.4, 1H), 7.02 (s, 1H), 4.01 (s, 3H), 3.92 (s, 3H).

Example 48***6-Chloro-N-(2-methoxycarbonyl-4,5-dimethoxyphenyl)-3-pyridinecarboxamide***

20 The title compound was prepared from 6-chloronicotinoyl chloride and 2-amino-4,5-dimethoxybenzoic acid methyl ester as a light yellow solid as described in Example 1. ¹H NMR (CDCl₃): 12.34 (s, 1H), 9.09 (d, J = 2.4, 1H), 8.62 (s, 1H), 8.29-8.26 (m, 1H), 7.52 (s, 1H), 7.50 (d, J = 8.4, 1H), 4.03 (s, 3H), 3.96 (s, 3H), 3.93 (s, 3H).

Example 49***6-Chloro-N-(2-methylcarbonyl-4,5-methylenedioxyphenyl)-3-pyridinecarboxamide***

25 The title compound was prepared from 6-chloronicotinoyl chloride and 2-amino-4,5-methylenedioxyacetophenone as a light grey-yellow solid as described in Example 1.

30

¹H NMR (CDCl₃): 9.09-9.08 (m, 1H), 8.55 (s, 1H), 8.30-8.27 (m, 1H), 7.50-7.47 (m, 1H), 7.35 (s, 1H), 6.10 (s, 2H), 2.64 (s, 3H).

Example 50

6-(2,2,2-Trifluoroethoxy)-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide

The title compound was prepared from 6-(2,2,2-trifluoroethoxy)pyridine-3-carboxylic acid and 4-ethoxy-2-nitroaniline as a yellow solid as described in Example 15.

¹H NMR (CDCl₃): 11.05 (s, 1H), 8.84-8.80 (m, 2H), 8.25-8.21 (m, 1H), 7.73 (d, J = 3.0, 1H), 7.32-7.28 (m, 1H), 7.01 (d, J = 8.7, 1H), 4.86 (q, J = 8.4, 2H), 4.11 (q, J = 6.9, 2H), 1.46 (t, J = 6.9, 3H).

Example 51

6-Chloro-N-(2,3-dinitro-4-methoxyphenyl)-3-pyridinecarboxamide

The title compound was prepared from 6-chloronicotinoyl chloride and 2,3-dinitro-4-methoxyaniline as a white solid as described in Example 1. ¹H NMR (CDCl₃): 9.99 (s, 1H), 8.96 (d, J = 2.4, 1H), 8.77 (d, J = 9.3, 1H), 8.18-8.14 (m, 1H), 7.53 (d, J = 9.0, 1H), 7.45 (d, J = 9.3, 1H), 4.02 (s, 3H).

Example 52

6-Chloro-N-(2-nitro-4,6-dimethylphenyl)-3-pyridinecarboxamide

The title compound was prepared from 6-chloronicotinoyl chloride and 2-nitro-4,6-dimethylaniline as a light yellow solid as described in Example 1. ¹H NMR (CDCl₃): 9.09 (s, 1H), 8.97 (d, J = 2.4, 1H), 8.22-8.18 (m, 1H), 7.60 (d, J = 1.2, 1H), 7.50 (d, J = 8.1, 1H), 7.42 (s, 1H), 2.43 (s, 3H), 2.35 (s, 3H).

Example 53***4-Chloromethyl-N-(4-ethoxy-2-nitrophenyl)-benzoylamide***

The title compound was prepared from 4-chloromethylbenzoyl chloride and 4-ethoxy-2-nitroaniline as a yellow solid as described in Example 1. ¹H NMR (CDCl₃): 11.11 (s, 1H), 8.87 (d, J = 9.3, 1H), 7.98 (d, J = 8.4, 2H), 7.73 (d, J = 3.0, 1H), 7.56 (d, J = 8.4, 2H), 7.32-7.28 (m, 1H), 4.65 (s, 2H), 4.11 (q, J = 6.9, 2H), 1.46 (t, J = 6.9, 3H).

Example 54***6-Chloro-N-(4,5-dimethoxy-2-nitrophenyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinoyl chloride and 4,5-dimethoxy-2-nitroaniline as a yellow solid as described in Example 1. ¹H NMR (CDCl₃): 11.84 (s, 1H), 9.04 (d, J = 1.8, 1H), 8.66 (s, 1H), 8.25-8.21 (m, 1H), 7.77 (s, 1H), 7.53 (d, J = 8.7, 1H), 4.07 (s, 3H), 3.97 (s, 3H).

Example 55***6-Chloro-N-(2-methoxy-5-pyridyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinoyl chloride and 5-amino-2-methoxypyridine as a pink solid as described in Example 1. ¹H NMR (CDCl₃): 8.87 (d, J = 2.1, 1H), 8.28 (d, J = 3.0, 1H), 8.20-8.17 (m, 1H), 7.99-7.96 (m, 1H), 7.67 (s, 1H), 7.49 (d, J = 8.7, 1H), 6.81 (d, J = 8.7, 1H), 3.95 (s, 3H).

Example 56***6-Chloro-N-(2-chloro-5-pyridyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinoyl chloride and 5-amino-2-chloropyridine as a white solid as described in Example 1. ¹H NMR (CDCl₃): 8.89-8.84

(m, 1H), 8.52 (d, J = 2.1, 1H), 8.27-8.23 (m, 1H), 8.21-8.17 (m, 1H), 7.83 (s, 1H), 7.53-7.50 (m, 1H), 7.39 (d, J = 8.7, 1H).

Example 57

5

6-Chloro-N-(4,6-dichloro-5-pyrimidyl)-3-pyridinecarboxamide

The title compound was prepared from 6-chloronicotinoyl chloride and 5-amino-4,6-dichloropyrimidine as a white solid as described in Example 1. ¹H NMR (CDCl₃): 8.95 (d, J = 2.4, 1H), 8.78 (s, 1H), 8.25-8.22 (m, 1H), 7.57 (s, 1H), 7.54 (d, J = 8.4, 1H).

10

Example 58

6-Cyano-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide

The title compound was prepared from 6-cyanonicotinic acid and 4-ethoxy-2-nitroaniline as a yellow solid as described in Example 15. ¹H NMR (CDCl₃): 11.22 (s, 1H), 9.31-9.30 (m, 1H), 8.81 (d, J = 9.3, 1H), 8.42-8.38 (m, 1H), 7.90-7.87 (m, 1H), 7.76 (d, J = 3.0, 1H), 7.34-7.30 (m, 1H), 4.12 (q, J = 6.9, 2H), 1.47 (t, J = 6.9, 3H).

15

Example 59

2-Methyl-N-(4-ethoxy-2-nitrophenyl)-3-furancarboxamide

The title compound was prepared from 2-methyl-3-furoic acid and 4-ethoxy-2-nitroaniline as an orange yellow solid as described in Example 15. ¹H NMR (CDCl₃): 10.66 (s, 1H), 8.82 (d, J = 9.3, 1H), 7.70 (d, J = 3.0, 1H), 7.34 (d, J = 1.8, 1H), 7.28-7.24 (m, 1H), 6.68 (d, J = 1.8, 1H), 4.09 (q, J = 6.9, 2H), 2.67 (s, 3H), 1.45 (t, J = 6.9, 3H).

25

Example 60***6-Chloro-N-(4-chloro-2-trifluoromethylphenyl)-3-pyridinecarboxamide***

The title compound was prepared from 6-chloronicotinoyl chloride and 4-chloro-2-trifluoromethylaniline as a white solid as described in Example 1. ¹H NMR (CDCl₃): 8.87 (d, J = 2.7, 1H), 8.32 (d, J = 8.7, 1H), 8.16-8.12 (m, 1H), 8.06 (bs, 1H), 7.82 (d, J = 2.4, 1H), 7.53-7.50 (m, 1H), 7.12 (d, J = 8.7, 1H).

Example 61***4-Chloro-2-nitro-N-(6-chloro-3-pyridyl)-benzoylamide***

The title compound was prepared from 4-chloro-2-nitrobenzoic acid and 5-amino-2-chloropyridine as a white solid as described in Example 15.

Example 62***6-Chloro-N-(2-pyrazinyl)-3-pyridinecarboxamide***

To a mixture of 6-chloronicotinoyl chloride (160 mg, 0.91 mmol), aminopyrazine (87 mg, 0.91 mmol) and THF (10 ml) was added triethylamine (0.25 ml, 1.82 mmol) and the mixture was stirred overnight at room temperature. The mixture was diluted with ethyl acetate (100 ml), wash with water (50 ml), saturated aqueous NaCl (50 ml), dried over anhydrous sodium sulfate and evaporated. The crude product was purified by column chromatography using hexane:ethyl acetate (3:2) to give 8 mg (4% yield) of the title compound as yellow solid. ¹H NMR (CDCl₃): 9.68 (d, J = 1.2 Hz, 1H), 8.95 (d, J = 2.7 Hz, 1H), 8.45 (d, J = 2.7 Hz, 1H), 8.4 (bs, 1H), 8.33- 8.32 (m, 1H), 8.23 (dd, J = 2.4, 8.1 Hz, 1H), 7.53 (dd, J = 0.6, 8.4 Hz, 1H).

Example 63***4-Chloro-2-nitro-N-(6-methoxy-3-pyridyl)-benzoylamide***

A mixture of cyanuric chloride (183 mg, 0.99 mmol) and 4-chloro-2-nitrobenzoic acid (200 mg, 0.99 mmol) in THF (5 ml) was stirred at room temperature for 30 minutes. To the mixture was added a solution of 5-amino-2-methoxy-pyridine (123 mg, 0.99 mmol) in THF (5 ml), followed by triethylamine (0.28 ml, 1.98 mmol). The mixture was stirred overnight at room temperature. It was diluted with 100 ml of hexane:ethyl acetate (1:1), wash with 2N NaOH (50 ml), water (50 ml), followed by saturated aqueous NaCl (50 ml), dried over anhydrous sodium sulfate and evaporated. The crude product was purified by column chromatography using hexane:ethyl acetate (3:1) to give 16 mg (5% yield) of the title compound as a pink solid. ¹H NMR (Acetone-d₆): 9.94 (s, 1H), 8.16 (d, *J* = 1.8 Hz, 1H), 8.07-7.86 (m, 4H), 6.81 (d, *J* = 9.3 Hz, 1H), 3.89 (s, 3H).

Example 64***6-Chloro-N-(2-cyano-6-chloromethyl-4-oxide-3-pyrazinyl)-3-pyridinecarboxamide***

The title compound was prepared from 3-amino-6-(chloromethyl)-2-pyrazine carbonitrile 4-oxide (203 mg, 1.1 mmol) and 6-chloronicotinoyl chloride (200 mg, 1.1 mmol) and was obtained as a yellow solid as described in Example 62. ¹H NMR (Acetone-d₆): 9.00 (s, 1H), 8.66 (s, 1H), 8.42 (dd, *J* = 8.1, 2.1 Hz, 1H), 7.65 (d, *J* = 8.1 Hz, 1H), 7.52 (s, 1H), 5.38 (s, 2H).

Example 65***6-Chloro-N-(2-cyano-4-methylphenyl)-3-pyridinecarboxamide***

A mixture of 2-amino-5-methyl-benzonitrile (80 mg, 0.61 mmol) and 6-chloronicotinoyl chloride (162 mg, 0.92 mmol) in anhydrous pyridine (5 ml) was refluxed for 4 hours. The mixture was diluted with 50 ml of hexane:ethyl acetate (1:1), wash with 2N HCl (25 ml), water (25 ml), followed by saturated aqueous NaCl (25 ml), dried over anhydrous sodium sulfate, and evaporated. The crude product was purified by column

chromatography using hexane:ethyl acetate (2:1) to give 30 mg (18% yield) of the title compound as pale pink solid. ¹H NMR (DMSO-d₆): 10.82 (s, 1H), 8.99 (d, *J* = 2.4 Hz, 1H), 8.38 (dd, *J* = 2.4, 8.1 Hz, 1H), 7.78-7.73 (m, 2H), 7.61-7.48 (m, 2H), 2.38 (s, 3H).

5

Example 66

6-Chloro-N-(4-chloro-6-methyl-2-nitrophenyl)-3-pyridinecarboxamide

The title compound was prepared from 4-chloro-2-methyl-6-nitroaniline (204 mg, 1.09 mmol) and 6-chloronicotinoyl chloride (192 mg, 1.09 mmol) and obtained as a light yellow solid as described in Example 62. ¹H NMR (CDCl₃): 9.13 (s, 1H), 8.96 (dd, *J* = 0.6, 2.4 Hz, 1H), 8.20 (dd, *J* = 2.7, 8.4 Hz, 1H), 7.93 (d, *J* = 2.4 Hz, 1H), 7.59 (d, *J* = 2.4 Hz, 1H), 7.51 (dd, *J* = 0.6, 8.1 Hz, 1H), 2.38 (s, 3H).

10

Example 67

4-Trifluoromethyl-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide

15

A mixture of 4-(trifluoromethyl)pyridine-3-carboxylic acid (100 mg, 0.52 mmol), 4-ethoxy-2-nitroaniline (94.7 mg, 0.52 mmol), and 1-(3-dimethylamino-propyl)-3-ethylcarbodiimide hydrochloride (199.3 mg, 1.04 mmol) in THF (5 ml) was refluxed overnight. The mixture was cooled to room temperature, diluted with 100 ml of hexane:ethyl acetate (1:1), wash with 2N NaOH (50 ml), water (50 ml), saturated aqueous NaCl (50 ml), dried over anhydrous sodium sulfate and evaporated. The crude product was purified by column chromatography using hexane:ethyl acetate (2:1) to give 17.3 mg (9.4% yield) of the title compound as yellow solid. ¹H NMR (CDCl₃): 10.48 (s, 1H), 9.01-8.96 (m, 2H), 8.74 (d, *J* = 9.3 Hz, 1H), 7.74-7.69 (m, 2H), 7.32 (dd, *J* = 3.0, 9.3 Hz, 1H), 4.13 (q, *J* = 6.9 Hz, 2H), 1.48 (t, *J* = 6.9 Hz, 3H).

20

25

Example 68***4-Bromomethyl-3-nitro-N-(6-chloro-3-pyridyl)-benzoylamide***

A mixture of 4-bromomethyl-3-nitrobenzoic acid (169 mg, 0.65 mmol), 5-amino-2-chloropyridine (83 mg, 0.65 mmol) and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (210 mg) in THF (10 ml) was refluxed for 24 h. The mixture was cooled to room temperature and diluted with hexane:ethyl acetate (1:1, 100 ml), washed with water (50 ml), 2N HCl (50 ml), brine (50 ml) and dried over anhydrous sodium sulfate. The solution was concentrated in vacuo to give a yellow solid which was washed with dichloromethane and dried to give 68 mg (28%) of the title compound. ¹H NMR (DMSO-d₆): 10.91 (s, 1H), 8.77 (d, J = 2.1 Hz, 1H), 8.63 (s, 1H), 8.31 (d, J = 8.1 Hz, 1H), 8.24 (dd, J = 2.4, 8.7 Hz, 1H), 7.96 (d, J = 8.1 Hz, 1H), 7.56 (d, J = 8.7 Hz, 1H), 5.11 (s, 2H).

Example 69***6-Chloromethyl-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide***

A) **Methyl 6-chloromethylnicotinate:** A mixture of methyl 6-methylnicotinate (2.3 g, 15.2 mmol) and N-chlorosuccinimide (4.6 g, 34 mmol) in chloroform was refluxed for 24 h to give 1.3 g of the title compound as a white solid.

B) **6-Chloromethylnicotinic acid:** A mixture of methyl 6-chloromethylnicotinate (1.3 g, 6.5 mmol) in 2N HCl (40 ml) was refluxed for 5 h. The mixture was cooled to room temperature to give 1.1 g of the title compound as a tan solid.

C) **6-Chloromethyl-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide:** A mixture of 6-chloromethylnicotinic acid (155 mg, 0.69 mmol), 4-ethoxy-2-nitroaniline (146 mg), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (238 mg) and pyridine in THF (10 ml) was refluxed for 17 h. The mixture was cooled to room temperature, diluted with hexane:ethyl acetate (1:1, 100 ml), washed with water (50 ml), 2N HCl (50 ml), brine (50 ml) and dried over anhydrous sodium sulfate. The solution was concentrated in vacuo and the product was purified by column chromatography using hexane:ethyl acetate (3:1) to give the title compound as a yellow solid (30 mg). ¹H NMR (CDCl₃): 11.14 (s, 1H), 9.18 (m, 1H), 8.84 (d, J = 9.3 Hz, 1H), 8.30 (dd, J = 2.4, 8.1 Hz,

1H), 7.75 (d, J = 3.0 Hz, 1H), 7.68 (d, J = 8.4 Hz, 1H), 7.31 (dd, J = 3.0, 9.3 Hz, 1H), 4.76 (s, 2H), 4.12 (m, 2H), 1.47 (t, J = 7.1 Hz, 3H).

Example 70

N-(4-Ethoxy-2-nitrophenyl)-3-furancarboxamide

The title compound was prepared from 3-furoic acid and 4-ethoxy-2-nitroaniline as a yellow solid as described in Example 15. ¹H NMR (DMSO-*d*₆): 10.25 (s, 1H), 8.36 (d, J=2.4, 1H), 7.83-7.81 (m, 1H), 7.53-7.47 (m, 2H), 7.35-7.31 (m, 1H), 6.94 (d, J=2.4, 1H), 4.12 (q, J=6.9, 2H), 1.35 (t, J=6.9, 3H).

Example 71

Identification of N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide and Analogs as Caspase Cascade Activators and Inducers of Apoptosis in Solid Tumor Cells

Human breast cancer cell lines T47D and ZR75-1 were grown according to media component mixtures designated by American Type Culture Collection + 10% FCS (Invitrogen Corporation, Life Technologies Division), in a 5% CO₂ -95% humidity incubator at 37 °C. T47D and ZR75-1 cells were maintained at a cell density between 30 and 80% confluency at a cell density of 0.1 to 0.6 x 10⁶ cells/ml. Cells were harvested at 600xg and resuspended at 0.65 x 10⁶ cells/ml into appropriate media + 10% FCS. An aliquot of 45 µl of cells was added to a well of a 96-well microtiter plate containing 5 µl of a 10% DMSO in RPMI-1640 media solution containing 1.6 to 100 µM of N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide or other test compound (0.16 to 10 µM final). An aliquot of 45 µl of cells was added to a well of a 96-well microtiter plate containing 5 µl of a 10% DMSO in RPMI-1640 media solution without test compound as the control sample. The samples were mixed by agitation and then incubated at 37 °C for 24 h in a 5% CO₂-95% humidity incubator. After incubation, the samples were removed from the incubator and 50 µl of a solution containing 20 µM of N-(Ac-DEVD)-N'-ethoxycarbonyl-R110 (SEQ ID NO:1) fluorogenic substrate (Cytovia, Inc.; WO99/18856), 20% sucrose (Sigma), 20 mM DTT (Sigma), 200 mM NaCl (Sigma), 40 mM Na PIPES buffer pH 7.2 (Sigma),

and 500 µg/ml lysolecithin (Calbiochem) was added. The samples were mixed by agitation and incubated at room temperature. Using a fluorescent plate reader (Model 1420 Wallac Instruments), an initial reading ($T = 0$) was made approximately 1- 2 min after addition of the substrate solution, employing excitation at 485 nm and emission at 530 nm, to determine the background fluorescence of the control sample. After the 3 h incubation, the samples were read for fluorescence as above ($T = 3$ h).

Calculation:

The Relative Fluorescence Unit values (RFU) were used to calculate the sample readings as follows:

$$\text{RFU}_{(T=3h)} - \text{Control RFU}_{(T=0)} = \text{Net RFU}_{(T=3h)}.$$

The activity of caspase cascade activation was determined by the ratio of the net RFU value for N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide or other test compound to that of control samples. The EC_{50} (nM) was determined by a sigmoidal dose-response calculation (Prism 2.0, GraphPad Software Inc.). The caspase activity (Ratio) and potency (EC_{50}) are summarized in Table I:

Table I. Caspase Activity and Potency

Example #	T47D		ZR75-1	
	Ratio	EC50 (nM)	Ratio	EC50 (nM)
1	8.9	2970	3.6	729
2	7.8	587	5.8	329
3	Inactive	Inactive	2.1	4620
4	Inactive	Inactive	3.2	4030
5	Inactive	Inactive	Inactive	Inactive
6	Inactive	Inactive	Inactive	Inactive
7	8.1	593	2.7	230
8	5.0	153	6.3	77
9	5.4	920	5.0	488
10	Inactive	Inactive	Inactive	Inactive
11	Inactive	Inactive	Inactive	Inactive
12	Inactive	Inactive	Inactive	Inactive
13	Inactive	Inactive	Inactive	Inactive
14	Inactive	Inactive	Inactive	Inactive
15	Inactive	Inactive	Inactive	Inactive
16	Inactive	Inactive	Inactive	Inactive
17	Inactive	Inactive	Inactive	Inactive
18	3.0	6750	5.1	3840
19	2.7	3550	3.8	6880
20	Inactive	Inactive	Inactive	Inactive
21	Inactive	Inactive	2.3	4670

22	7.6	466	6.6	295
23	7.2	608	9.9	171
24	Inactive	Inactive	Inactive	Inactive
25	5.8	345	6.6	133
26	5.5	42	3.2	20
27	Inactive	Inactive	Inactive	Inactive
28	2.4	7330	5.1	4510
29	5.8	1800	6.4	1000
30	Inactive	Inactive	Inactive	Inactive

Thus, N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide (Example 29) and analogs are identified as potent caspase cascade activators and inducer of apoptosis in solid tumor cells.

5

Example 72

N-(4-Methoxy-2-nitrophenyl)-3-pyridinecarboxamide Induces Nuclear Fragmentation in Jurkat Cells

10

15

The ability of N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide to induce nuclear fragmentation was tested by treatment of Jurkat cells with the test compound followed by staining of the nucleus with Syto16, a fluorescent DNA dye (Molecular Probes, Eugene, OR). The nuclei of Jurkat cells treated with vehicle control (DMSO) are seen to be round with dispersed chromatin that is moderately stained with Syto16 (Fig. 1A). In contrast, Jurkat cells treated with 5 μ M of N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide for 24 h have shrunken and fragmented nuclei (Fig. 1B), which is a hallmark of caspase-mediated apoptosis. These results corroborate the caspase induction assays by showing that N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide can induce a key cellular marker of apoptosis.

Example 73

N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide Induces Mitotic Arrest in Jurkat Cells

Jurkat cells were incubated with a range of concentrations of N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide (0.02 μ M to 5 μ M) for 6 h under normal growth conditions; control cultures were treated with DMSO vehicle. The cells were then treated for 20 min. with 800 nM Syto 16 (Molecular Probes, Eugene, OR). Cytospin preparations were then prepared and the samples were viewed by fluorescent microscopy using a fluorescein filter set. For each concentration of test compound, the number of mitotic figures were counted and expressed as a percentage of the total number of cells. Three fields from each condition were evaluated and the mean and SEM were calculated and plotted as a function of drug concentration (Fig. 2). The results show that N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide is an effective inducer of mitotic arrest in Jurkat cells up to a concentration of 5 μ M.

Example 74

Treatment with N-(4-Methoxy-2-nitrophenyl)-3-pyridinecarboxamide or 6-Chloro-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide Leads to Cell Cycle Arrest and Apoptosis in Solid Tumor Cell Line

T47D, a breast cancer cell line, was maintained and harvested as described in Example 71. 1×10^6 cells were treated with 10 μ M of N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide or 6-chloro-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide for 48 h at 37°C. As a control, cells were also incubated with equivalent amount of solvent as control (DMSO). Cells were harvested at 1,200 rpm and washed twice with 5 mM EDTA/PBS. Cells were then resuspended in 300 μ l EDTA/PBS and 700 μ l of 100% ethanol, vortexed and incubated at room temperature for 1 h. Samples were spun down at 1,200 rpm for 5 min and the supernatant was removed. A solution containing 100 μ g/ml of propidium iodide and 1 mg/ml of RNase A (fresh) was added to the samples and incubated for 1 h at room temperature. Samples were then transferred to 12x75 mm polystyrene tubes and analyzed on a flow cytometer. All flow cytometry analyses were

performed on FACScalibur (Becton Dickinson) using Cell Quest analysis software. On the x-axis is plotted the fluorescence intensity and on the y-axis is plotted the number of cells with that fluorescence intensity. The T47D control cell population profile is seen in Fig. 3A and the increase in G2/M DNA content (M4) cells that is seen when treated with N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide at 10 μ M is seen in Fig. 3B. An increase in the sub-diploid DNA content of cells (marker M1 region, Fig 3) is also seen to increase from 2% to 25% with compound treatment. The sub-diploid amount of DNA (M1) is indicative of apoptotic cells which have undergone DNA degradation or fragmentation. In cells treated with 6-chloro-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarbox-amide, a more potent analog, at 1 μ M (Fig. 4B), the accumulation of T47D cells in the G2/M phase (M4) is similarly seen. In addition, the sub-G1 population of cells with reduced DNA content (M1) indicative of apoptosis is seen to increase substantially over control cells (Fig. 4A), indicating cell cycle arrest and induction of apoptosis by treatment with N-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide and its more potent analog 6-chloro-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide.

Example 75

6-Chloro-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide Inhibits the Proliferation of Solid Tumor HeLa Cell Line

HeLa cells were grown according to media component mixtures designated by American Type Culture Collection + 10% FCS (Invitrogen Corporation, Life Technologies Division), in a 5% CO₂ -95% humidity incubator at 37 °C. In a well of a 96 well plate, 30,000 cells were seeded and treated with compound at the indicated concentrations for 48 hr in a 5% CO₂- 95% humidity incubator at 37°C. Control wells were treated with the same amount of solvent (DMSO) as the compound samples. After the indicated treatment time, the supernatant was removed to a sterile culture tube and the wells washed with phosphate buffered saline, and the adherent cells trypsinized for 5 min. The trypsinized cells were added to the culture supernatant, cells were collected (1,200 rpm, 10 min), washed with phosphate buffered saline, and resuspended in fresh media. The cells were counted for trypan blue negative cells, and the cells were diluted with fresh


media to 1,000 cells/ml. To a well of a 24-well plate, 0.1 ml of the cell suspension was added along with 1 ml of fresh media (cell suspensions were passed through a 22G needle several times just before plating to form single cell suspensions). Plates are incubated in a 5% CO₂- 95% humidity incubator at 37°C for 5-7 days. Colonies are counted when the sizes reached greater than 50 cells per colony. Cells are washed with phosphate buffered saline, fixed with 100% methanol for 15 min, and then stained with 0.5% gentian violet for 15 min. Colonies are rinsed with water and the colonies counted and the fraction surviving expressed as the percentage of the number of control colonies.

The results showed that after a 48 hr treatment, 6-chloro-N-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide inhibits the ability of HeLa cells to continue to proliferate and inhibited their colony forming ability with an IC₅₀ of about 100 nM (Fig. 5)

Having now fully described this invention, it will be understood by those of ordinary skill in the art that the same may be performed within a wide and equivalent range of conditions, formulations and other parameters without affecting the scope of the invention or any embodiment thereof. All patents, patent applications and publications cited herein are fully incorporated by reference herein in their entirety.

What Is Claimed Is:

1. A method of treating a disorder responsive to the induction of apoptosis in an animal suffering therefrom, comprising administering to a mammal in need of such treatment an effective amount of a compound of Formula V:




or a pharmaceutically acceptable salt or prodrug thereof, wherein:

Ar' and Ar are independently optionally substituted aryl or optionally substituted heteroaryl, provided that the ring structure of said optionally substituted heteroaryl comprises not more than two nitrogen atoms; and

R₁₁ is hydrogen; or alkyl, cycloalkyl, aryl or heteroaryl, each of which is optionally substituted.

2. The method of claim 1, wherein said compound is of Formula I:



or pharmaceutically acceptable salts or prodrugs thereof, wherein:

A is N or C-R₆, B is N or C-R₉, D is N or C-R₁₀, E is N or C-R₆ and F is N or C-R₇, provided that not more than two of A, B, D, E and F are N in the same time;

Ar is optionally substituted and is an aryl or heteroaryl;

R₆-R₁₀ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl,

heterocycloalkyl, hydroxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido or alkylthiol; and

R₁₁ is hydrogen or optionally substituted alkyl, cycloalkyl, aryl, or heteroaryl.

- 5 3. The method of claim 2, wherein A is N, B is C-R₉ and F is C-R₇.
4. The method of claim 1, wherein Ar' is optionally substituted and is furyl, pyrrolyl, pyrazolyl, imidazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl or phenyl.
- 10 5. The method of claim 4, wherein Ar' is optionally substituted furyl.
6. The method of claim 5, wherein Ar is optionally substituted phenyl.
7. The method of claim 6, wherein Ar' is optionally substituted 3-furyl.
- 15 8. The method of claim 7, wherein Ar' is unsubstituted 3-furyl.
9. The method of claim 8, wherein said compound is *N*-(4-ethoxy-2-nitrophenyl)-3-furancarboxamide.
- 20 10. The method of claim 4, wherein Ar' is optionally substituted pyrrolyl.
11. The method of claim 10, wherein Ar is optionally substituted phenyl.
- 25 12. The method of claim 11, wherein Ar' is optionally substituted 3-pyrrolyl.
13. The method of claim 12, wherein Ar' is unsubstituted 3-pyrrolyl.
- 30 14. The method of claim 13, wherein said compound is *N*-(4-ethoxy-2-nitrophenyl)-3-pyrrolecarboxamide.

15. The method of claim 4, wherein Ar' is optionally substituted phenyl.

16. The method of claim 15, wherein Ar is optionally substituted phenyl.

5 17. The method of claim 16, wherein said compound is selected from the group consisting of:

4-Chloro-*N*-(4-ethoxy-2-nitrophenyl)-benzoylamide; and

4-Chloromethyl-*N*-(4-ethoxy-2-nitrophenyl)-benzoylamide.

10 18. The method of claim 15, wherein Ar is optionally substituted pyridyl.

19. The method of claim 18, wherein said compound is selected from the group consisting of:

4-Chloro-2-nitro-*N*-(6-chloro-3-pyridyl)-benzoylamide;

15 4-Chloro-2-nitro-*N*-(6-methyl-3-pyridyl)-benzoylamide; and

4-Bromomethyl-3-nitro-*N*-(6-chloro-3-pyridyl)-benzoylamide.

20. The method of claim 4, wherein Ar' is optionally substituted pyrazinyl.

20 21. The method of claim 20, wherein Ar is optionally substituted phenyl.

22. The method of claim 21, wherein Ar' is 3-pyrazinyl.

25 23. The method of claim 22, wherein said compound is selected from the group consisting of:

6-Methyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyrazinecarboxamide; and

N-(4-Ethoxy-2-nitrophenyl)-3-pyrazinecarboxamide.

30 24. The method of claim 4, wherein Ar' is optionally substituted pyrimidinyl.

25. The method of claim 24, wherein Ar is optionally substituted phenyl.

26. The method of claim 25, wherein Ar' is optionally substituted 5-pyrimidinyl.

27. The method of claim 26, wherein said compound is *N*-(4-ethoxy-2-nitrophenyl)-5-pyrimidinecarboxamide.

28. The method of claim 4, wherein Ar' is optionally substituted pyridyl.

29. The method of claim 28, wherein Ar is optionally substituted phenyl.

30. The method of claim 29, wherein Ar' is optionally substituted 2-pyridyl.


31. The method of claim 30, wherein said compound is selected from the group consisting of:

N-(4-Ethoxy-2-nitrophenyl)-2-pyridinecarboxamide; and

N-(4-Ethoxy-2-nitrophenyl)-1-*N*-oxide-2-pyridinecarboxamide.

32. The method of claim 29, wherein Ar' is optionally substituted 3-pyridyl.

33. The method of claim 32, wherein said compound is of Formula III:



or a pharmaceutically acceptable salt or prodrug thereof, wherein

R₁-R₇ and R₉-R₁₀ are independently hydrogen, halo, haloalkyl, haloalkoxy, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl,

arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, aminoalkyl, cyano, cyanoalkyl, acyl, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, alkoxycarbonyl, aryloxy, arylalkoxy, carboxy, carbonylamido or alkylthiol; and

5 R_{11} is hydrogen; or alkyl, cycloalkyl, aryl or heteroaryl, each of which is optionally substituted.

34. The method of claim 33, wherein R_1 and R_2 , or R_2 and R_3 , or R_3 and R_4 , or R_4 and R_5 are taken together to form an optionally substituted carbocycle or an optionally substituted heterocycle.

35. The method of claim 34, wherein said R_1 and R_2 , or R_2 and R_3 , or R_3 and R_4 , or R_4 and R_5 are taken together to form $-OCH_2O-$, $-(CH_2)_3-$, $-(CH_2)_4-$, $-OCH_2CH_2O-$, $-CH_2N(R)CH_2-$, $-CH_2CH_2N(R)CH_2-$, $-CH_2N(R)CH_2CH_2-$, $-CH=CH-CH=CH-$, $-N(R)-CH=CH-$, $-CH=CH-N(R)-$, $-O-CH=CH-$, $-CH=CH-O-$, $-S-CH=CH-$, $-CH=CH-S-$, $-N=CH-CH=CH-$, $-CH=N-CH=CH-$, $-CH=CH-N=CH-$, $-CH=CH-CH=N-$ or $-N=CH-CH=N-$, wherein the carbocycle or heterocycle is optionally substituted, and R is hydrogen, alkyl, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl or aminoalkyl.

36. The method of claim 33, wherein R_6 , R_7 and R_{10} are independently hydrogen or fluoro.

37. The method of claim 33, wherein R_1 is nitro.


38. The method of claim 33, wherein R_2 , R_4 , and R_5 are independently hydrogen or fluoro.

-67-

39. The method of claim 33, wherein said compound is selected from the group consisting of:

N-(4-Methyl-2-nitrophenyl)-3-pyridinecarboxamide;
N-(4-Ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
 5 *N*-(4-Methoxy-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4,5-difluoro-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(3-bromo-4-methoxy-6-nitrophenyl)-3-pyridinecarboxamide;
 5,6-Dichloro-*N*-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(2-methyl-4-methoxyphenyl)-3-pyridinecarboxamide;
 10 6-Chloro-*N*-(4-ethoxy-2-nitrophenyl)-*N*-methyl-3-pyridinecarboxamide;
 6-Chloro-*N*-(2-cyano-4,5-dimethoxyphenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-chloro-2-trifluoromethylphenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-chloro-2-cyanophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(2,4-dimethyl-6-nitrophenyl)-3-pyridinecarboxamide;
 15 6-Chloro-*N*-(3,4-dimethoxy-6-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(2-cyano-4-methylphenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-chloro-2-methyl-6-nitrophenyl)-3-pyridinecarboxamide; and
 4-Trifluoromethyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide.

40. The method of claim 33, wherein said compound is of Formula IV:




or a pharmaceutically acceptable salt or prodrug thereof.

41. The method of claim 40, wherein said compound is selected from the group consisting of:

6-Chloro-*N*-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide;

6-Chloro-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-methyl-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-methoxy-2-nitrophenyl)-1-*N*-oxide-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-chloro-2-nitrophenyl)-3-pyridinecarboxamide;
 5 6-Fluoro-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-fluoro-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-trifluoromethyl-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(2-nitro-4-trifluoromethoxyphenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-benzyloxy-2-nitrophenyl)-3-pyridinecarboxamide;
 10 6-Methyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-cyano-2-nitrophenyl)-3-pyridinecarboxamide;
 6-(2,2,2-Trifluoroethoxy)-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Dimethylamino-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-*t*-butyl-2-nitrophenyl)-3-pyridinecarboxamide;
 15 6-Trifluoromethyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide; and
 6-Chloromethyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide.

42. A method for treating or preventing cancer, comprising administering to an
 animal in need of such treatment an effective amount of a compound of Formula V:




or a pharmaceutically acceptable salt or prodrug thereof, wherein:

25 Ar' and Ar are independently optionally substituted aryl or optionally substituted
 heteroaryl, provided that the ring structure of said optionally substituted heteroaryl
 comprises not more than two nitrogen atoms; and

R₁₁ is hydrogen; or alkyl, cycloalkyl, aryl or heteroaryl, each of which is optionally substituted.

43. The method of claim 42, wherein said compound is of Formula I:



5 or pharmaceutically acceptable salts or prodrugs thereof, wherein:

A is N or C-R₈, B is N or C-R₉, D is N or C-R₁₀, E is N or C-R₆ and F is N or C-R₇, provided that not more than two of A, B, D, E and F are N in the same time;

Ar is optionally substituted and is an aryl or heteroaryl;

10 R₆-R₁₀ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido or alkylthiol; and

R₁₁ is hydrogen or optionally substituted alkyl, cycloalkyl, aryl, or heteroaryl.

15

44. The method of claim 42, wherein Ar' is optionally substituted and is furyl, pyrrolyl, pyrazolyl, imidazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl or phenyl.

20 45. The method of claim 42, wherein said cancer is selected from the group consisting of Hodgkin's disease, non-Hodgkin's lymphoma, acute lymphocytic leukemia, chronic lymphocytic leukemia, multiple myeloma, neuroblastoma, breast carcinoma, ovarian carcinoma, lung carcinoma, Wilms' tumor, cervical carcinoma, testicular carcinoma, soft-tissue sarcoma, primary macroglobulinemia, bladder carcinoma, chronic granulocytic leukemia, primary brain carcinoma, malignant melanoma, small-cell lung carcinoma, stomach carcinoma, colon carcinoma, malignant pancreatic insulinoma, malignant carcinoid carcinoma, malignant melanoma, choriocarcinoma, mycosis fungoides, head or neck carcinoma, osteogenic sarcoma, pancreatic carcinoma, acute granulocytic leukemia, hairy cell leukemia, neuroblastoma, rhabdomyosarcoma, Kaposi's


25

-70-

sarcoma, genitourinary carcinoma, thyroid carcinoma, esophageal carcinoma, malignant hypercalcemia, cervical hyperplasia, renal cell carcinoma, endometrial carcinoma, polycythemia vera, essential thrombocytosis, adrenal cortex carcinoma, skin cancer and prostatic carcinoma.

5

46. A method for the treatment of drug resistant cancer, comprising administering to an animal in need of such treatment an effective amount of a compound of the Formula V:



10

or a pharmaceutically acceptable salt or prodrug thereof, wherein:


Ar' and Ar are independently optionally substituted aryl or optionally substituted heteroaryl, provided that the ring structure of said optionally substituted heteroaryl comprises not more than two nitrogen atoms; and

15

R₁₁ is hydrogen; or alkyl, cycloalkyl, aryl or heteroaryl, each of which is optionally substituted.

47. The method of claim 46, wherein said compound is of Formula I:

20



or pharmaceutically acceptable salts or prodrugs thereof, wherein:

A is N or C-R₈, B is N or C-R₉, D is N or C-R₁₀, E is N or C-R₆ and F is N or C-R₇, provided that not more than two of A, B, D, E and F are N in the same time;

25

Ar is optionally substituted and is an aryl or heteroaryl;

R₆-R₁₀ are independently hydrogen, halo, haloalkyl, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, cyano, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, carboxy, carbonylamido or alkylthiol; and

R₁₁ is hydrogen or optionally substituted alkyl, cycloalkyl, aryl, or heteroaryl.

48. The method of claim 46, wherein Ar' is optionally substituted and is furyl, pyrrolyl, pyrazolyl, imidazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl or phenyl.

49. The method of claim 42 or 46, additionally comprising administering at least one known cancer chemotherapeutic agent, or a pharmaceutically acceptable salt of said agent.

50. The method of claim 49, wherein said known cancer therapeutic agent is selected from the group consisting of busulfan, cis-platin, mitomycin C, carboplatin, colchicine, vinblastine, paclitaxel, docetaxel, camptothecin, topotecan, doxorubicin, etoposide, 5-azacytidine, 5-fluorouracil, methotrexate, 5-fluoro-2'-deoxy-uridine, ara-C, hydroxyurea, thioguanine, melphalan, chlorambucil, cyclophosphamide, ifosfamide, vincristine, mitoguanzone, epirubicin, aclarubicin, bleomycin, mitoxantrone, elliptinium, fludarabine, octreotide, retinoic acid, tamoxifen, Herceptin®, Rituxan® and alanosine.


51. The method of claim 42 or 46, additionally comprising treating said animal with radiation-therapy.

52. The method of claim 42 or 46, wherein said compound is administered after the surgical treatment of said animal for cancer.

53. The method of claim 1, wherein said disorder is an autoimmune disease.

54. The method of claim 1, wherein said disorder is rheumatoid arthritis.

55. The method of claim 1, wherein said disorder is inflammatory bowel disease.
56. The method of claim 1, wherein said disorder is a skin disease.
57. The method of claim 56, wherein said disorder is psoriasis.
58. A compound of Formula III:



or a pharmaceutically acceptable salt or prodrug thereof, wherein:

R_1 and R_5 are independently selected from the group consisting of hydrogen, hydroxy, alkyl, alkoxy, halogen, NO_2 , cyano, haloalkyl, haloalkoxy, amino and aminoalkyl, provided that at least one of R_1 and R_5 is selected from the group consisting of NO_2 , cyano, alkyl and haloalkyl;

R_2 and R_4 are independently selected from the group consisting of hydrogen, hydroxy, halogen, cyano, haloalkyl, haloalkoxy, amino and aminoalkyl;

R_3 is alkyl, Cl, F, haloalkyl, alkoxy, arylalkoxy, cyano, haloalkyloxy, amino or aminoalkyl;

R_6 is hydrogen, hydroxy, alkyl, NO_2 , cyano, haloalkyl, haloalkyloxy, amino or aminoalkyl;

R_7 is hydrogen, hydroxy, alkyl, NO_2 , cyano, haloalkyl, haloalkyloxy, amino or aminoalkyl;

R_9 is hydroxy, alkyl, halogen, NO_2 , haloalkyl, alkoxy, cyano, haloalkyloxy, amino or aminoalkyl;


R₁₀ is hydrogen, hydroxy, alkyl, Cl, F, NO₂, cyano, haloalkyl, haloalkyloxy, amino or aminoalkyl; and

R₁₁ is hydrogen, alkyl or haloalkyl.

5 59. The compound of claim 58, wherein said compound is selected from the group consisting of:

6-Chloro-*N*-(4,5-difluoro-2-nitrophenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(3-bromo-4-methoxy-6-nitrophenyl)-3-pyridinecarboxamide;
5,6-Dichloro-*N*-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide;
10 6-Chloro-*N*-(2-methyl-4-methoxyphenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(4-ethoxy-2-nitrophenyl)-*N*-methyl-3-pyridinecarboxamide;
6-Chloro-*N*-(2-cyano-4,5-dimethoxyphenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(4-chloro-2-trifluoromethylphenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(4-chloro-2-cyanophenyl)-3-pyridinecarboxamide;
15 6-Chloro-*N*-(2,4-dimethyl-6-nitrophenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(3,4-dimethoxy-6-nitrophenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(2-cyano-4-methylphenyl)-3-pyridinecarboxamide;
6-Chloro-*N*-(4-chloro-2-methyl-6-nitrophenyl)-3-pyridinecarboxamide; and
4-Trifluoromethyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide.

20 60. The compound of claim 58, wherein said compound is of Formula IV:




25 or a pharmaceutically acceptable salt or prodrug thereof.

61. The compound of claim 60, wherein said compound is selected from the group consisting of:

6-Chloro-*N*-(4-methoxy-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-methyl-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-methoxy-2-nitrophenyl)-1-*N*-oxide-3-pyridinecarboxamide;
 5 6-Chloro-*N*-(4-chloro-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Fluoro-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-fluoro-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-trifluoromethyl-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(2-nitro-4-trifluoromethoxyphenyl)-3-pyridinecarboxamide;
 10 6-Chloro-*N*-(4-benzyloxy-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Methyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Chloro-*N*-(4-cyano-2-nitrophenyl)-3-pyridinecarboxamide;
 6-(2,2,2-Trifluoroethoxy)-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Dimethylamino-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide;
 15 6-Chloro-*N*-(4-*t*-butyl-2-nitrophenyl)-3-pyridinecarboxamide;
 6-Trifluoromethyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide; and
 6-Chloromethyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyridinecarboxamide.

62. A compound of Formula (VI):



or a pharmaceutically acceptable salt or prodrug thereof, wherein

R_1 - R_5 , R_7 and R_9 - R_{10} are independently hydrogen, halo, haloalkyl, haloalkoxy, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl,

carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, aminoalkyl, cyano, cyanoalkyl, acyl, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, alkoxycarbonyl, aryloxy, arylalkoxy, carboxy, carbonylamido or alkylthiol; and

R_{11} is hydrogen; or alkyl, cycloalkyl, aryl or heteroaryl, each of which is optionally substituted.

63. The compound of claim 62, or a pharmaceutically acceptable salt thereof, with the proviso that at least one of R_1 and R_5 is selected from the group consisting of NO_2 , cyano, alkyl and haloalkyl.

64. The compound of claim 62 or a pharmaceutically acceptable salt or prodrug thereof, wherein:

R_1 and R_5 are independently selected from the group consisting of hydrogen, hydroxy, alkyl, halogen, NO_2 , cyano, haloalkyl, haloalkoxy, amino and aminoalkyl;

R_2 and R_4 are independently selected from the group consisting of hydrogen, hydroxy, halogen, haloalkyl, haloalkoxy, amino and aminoalkyl;

R_3 is alkyl, Cl, F, haloalkyl, alkoxy, arylalkoxy, cyano, haloalkoxy, amino or aminoalkyl;

R_7 , R_9 and R_{10} are independently selected from the group consisting of hydrogen, hydroxy, alkyl, halogen, NO_2 , cyano, haloalkyl, alkoxy, haloalkoxy, amino and aminoalkyl; and


R_{11} is hydrogen, alkyl or haloalkyl.

65. The compound of claim 64, wherein said compound is selected from the group consisting of:

6-Methyl-*N*-(4-ethoxy-2-nitrophenyl)-3-pyrazinecarboxamide; and

N-(4-Ethoxy-2-nitrophenyl)-3-pyrazinecarboxamide.

66. A compound of Formula (VII):



or a pharmaceutically acceptable salt or prodrug thereof, wherein:

5 R_1 - R_3 , R_5 - R_{10} are independently hydrogen, halo, haloalkyl, haloalkoxy, aryl, fused aryl, carbocyclic, a heterocyclic group, a heteroaryl group, alkyl, alkenyl, alkynyl, arylalkyl, arylalkenyl, arylalkynyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, carbocycloalkyl, heterocycloalkyl, hydroxyalkyl, nitro, amino, aminoalkyl, cyano, cyanoalkyl, acyl, acylamido, hydroxy, thiol, acyloxy, azido, alkoxy, alkoxycarbonyl, 10 aryloxy, arylalkoxy, carboxy, carbonylamido or alkylthiol; and

R_{11} is hydrogen; or alkyl, cycloalkyl, aryl or heteroaryl, each of which is optionally substituted.

67. The compound of claim 66, or a pharmaceutically acceptable salt thereof, 15 with the proviso that at least one of R_6 and R_7 is selected from the group consisting of NO_2 , cyano, alkyl and haloalkyl.

68. The compound of claim 67 or a pharmaceutically acceptable salt or prodrug thereof, wherein:

20 R_1 , R_2 , R_3 and R_5 are independently selected from the group consisting of hydrogen, hydroxy, alkyl, halogen, NO_2 , cyano, haloalkyl, alkoxy, haloalkoxy, amino and aminoalkyl;

R_6 and R_7 are independently selected from the group consisting of hydrogen, hydroxy, alkyl, halogen, NO_2 , cyano, haloalkyl, haloalkoxy, amino and aminoalkyl;

25 R_8 and R_{10} are independently selected from the group consisting of hydrogen, hydroxy, alkyl, halogen, NO_2 , cyano, haloalkyl, haloalkoxy, amino and aminoalkyl;

-77-

R₉ is hydroxy, alkyl, halogen, NO₂, cyano, haloalkyl, alkoxy, haloalkoxy, amino or aminoalkyl; and

R₁₁ is hydroxy, alkyl or haloalky.

5 69. The compound of claim 66, wherein said compound is selected from the group consisting of:

4-Chloro-2-nitro-*N*-(6-chloro-3-pyridyl)-benzoylamide;
4-Chloro-2-nitro-*N*-(6-methyl-3-pyridyl)-benzoylamide; and
4-Bromomethyl-3-nitro-*N*-(6-chloro-3-pyridyl)-benzoylamide.

10

70. A compound selected from the group consisting of:

N-(4-Ethoxy-2-nitrophenyl)-3-pyrrolylcarboxamide;
4-Chloro-*N*-(4-ethoxy-2-nitrophenyl)-benzoylamide;
4-Chloromethyl-*N*-(4-ethoxy-2-nitrophenyl)-benzoylamide;
15 *N*-(4-Ethoxy-2-nitrophenyl)-5-pyrimidinecarboxamide;
N-(4-Ethoxy-2-nitrophenyl)-2-pyridinecarboxamide; and
N-(4-Ethoxy-2-nitrophenyl)-1-*N*-oxide-2-pyridinecarboxamide.

15

20

71. A pharmaceutical composition, comprising the compound of any one of claims 58-70, and a pharmaceutically acceptable carrier.

25

72. The pharmaceutical composition of claim 71, further comprising at least one known cancer chemotherapeutic agent, or a pharmaceutically acceptable salt of said agent.

30

73. The pharmaceutical composition of claim 72, wherein said known cancer chemotherapeutic agent is selected from the group consisting of busulfan, cis-platin, mitomycin C, carboplatin, colchicine, vinblastine, paclitaxel, docetaxel, camptothecin, topotecan, doxorubicin, etoposide, 5-azacytidine, 5-fluorouracil, methotrexate, 5-fluoro-2'-deoxy-uridine, ara-C, hydroxyurea, thioguanine, melphalan, chlorambucil, cyclophosphamide, ifosfamide, vincristine, mitoguazone, epirubicin, aclarubicin, bleomycin,

mitoxantrone, elliptinium, fludarabine, octreotide, retinoic acid, tamoxifen, Herceptin®, Rituxan® and alanosine.

1/5




FIG.1A




FIG.1B

2/5






FIG.2

3/5

MARKER %

All	100.0
M1	2.6
M2	42.4
M3	3.8
M4	34.8
M5	9.8


FIG. 3A

MARKER %

All	100.0
M1	25.1
M2	7.4
M3	5.3
M4	57.2
M5	3.8


FIG. 3B

4/5



MARKER %	
All	100.0
M1	3.3
M2	68.4
M3	6.1
M4	16.5
M5	5.5

FIG. 4A



MARKER %	
All	100.0
M1	56.1
M2	13.7
M3	4.3
M4	18.5
M5	7.3

FIG. 4B

5/5




FIG.5

International Application No
PCT/US 01/02478

IPC 7	C07D213/82	C07D239/28	C07D241/28	C07D213/75	C07D401/12
	C07D307/68	A61K31/44	A61K31/4965	C07D207/34	A61K31/341

B. FIELDS SEARCHED

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

EPO-Internal, CHEM ABS Data, BEILSTEIN Data, PAJ, WPI Data

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 25723 A (AGREVO UK LTD ;RIORDAN PETER DOMINIC (GB); BODDY IAN KENNETH (NZ);) 28 September 1995 (1995-09-28) cited in the application the whole document ---	1-73
X	WO 99 36391 A (TULARIK INC) 22 July 1999 (1999-07-22) cited in the application see general formula and claims 41 and 64 ---	1-57
P,Y	US 6 028 111 A (CHAPLIN DAVID ET AL) 22 February 2000 (2000-02-22) the whole document ---	1-73
P,Y	WO 00 55114 A (CYTOVIA INC) 21 September 2000 (2000-09-21) the whole document ---	1-73

-/-

☒ Patent family members are listed in annex.

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- ***T** later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- ***X** document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- ***Y** document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- ***A** document member of the same patent family

21 May 2001

31.05.01.

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer _____

Scruton-Evans. I

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/02478

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO 01 21598 A (ASTRAZENECA UK LTD ;ASTRAZENECA AB (SE)) 29 March 2001 (2001-03-29) see examples 141,150 and definitions of R8 and claim 10	1,2
X	LIEGEOIS, JEAN FRANCOIS F. ET AL: "New pyridobenzodiazepine derivatives as potential antipsychotics: synthesis and neurochemical study" J. MED. CHEM. (1993), 36(15), 2107-14 , XP000986228 see Table II, compounds 21a-f	66,67
X	WHITE, G. A.: "Substituted 2-methylbenzanilides and structurally related carboxamides: inhibition of complex II activity in mitochondria from a wild-type strain and a carboxin-resistant mutant strain of Ustilago maydis" PESTIC. BIOCHEM. PHYSIOL. (1989), 34(3), 255-76 , XP000995985 see compound LXV, page 261	66,67
X	US 4 261 730 A (BOLLINGER FREDERIC G ET AL) 14 April 1981 (1981-04-14) see Table II	66
X	GB 1 573 576 A (MONSANTO CO) 28 August 1980 (1980-08-28) see general formula	66
X	ATWELL, G. J. ET AL: "Potential antitumor agents. VII. Bisquaternary salts" J. MED. CHEM. (1968), 11(2), 300-5 , XP000996549 see Table II	66,67
P,X	WO 01 02354 A (MORIMOTO MASAYUKI ;TOHNISHI MASANORI (JP); SEO AKIRA (JP); ENDOH K) 11 January 2001 (2001-01-11) see general formula and Q-27, pages 53->	66,67
X	COOPER, G. H. ET AL: "Reductive acylation of 5-nitropyridines with anhydrides of acetic, phthalic, and succinic acids" J. CHEM. SOC. C (1971), (19), 3257-60 , XP000996536 see compound 9, Scheme 2	66
X	US 3 406 168 A (SCHMIDT GUNTHER) 15 October 1968 (1968-10-15) see formula IV and examples	66,67
	-/--	

INTERNATIONAL SEARCH REPORT

 International Application No
 PCT/US 01/02478

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	JP 2000 256358 A (YAMANOUCHI PHARM CO LTD) 19 September 2000 (2000-09-19) see example 55	62
P,X	M.S.S.PALANKI: "Inhibitors of NF-B and Ap-1 Gene Expression" J MED CHEM, vol. 43, no. 21, 2000, pages 3995-4004, XP000986229 see Table 3, compound 103	62,71
P,X	WO 00 06550 A (ITO KUNIHITO ;MOCHIZUKI NOBUO (JP); NIPPON SODA CO (JP); UCHIDA SE) 10 February 2000 (2000-02-10) see compound 4-33	62,71
X	M BALCELLS: "Synthesis of Phenoxyphenyl Pyridine and Pyrazine carboxamides" J AGRIC FOOD CHEM, vol. 48, no. 1, 2000, pages 83-87, XP002154583 see compound 3d, Table 1	62
X	EP 0 807 633 A (PFIZER) 19 November 1997 (1997-11-19) See example 7 and preparations 9 and 10, page 28	62
X	M DOLEZAL: "Synthesis and photosynthesising Activity of some Anilides" CHEM PAPERS, vol. 53, no. 2, 1999, pages 126-130, XP000996425 see Table 1	62-64
P,X	M DOLEZAL: "Synthesis and Antimycobacterial, Antifungal and Photosynthesis-inhibiting evaluation of some Anilides" CHEM PAPERS, vol. 54, no. 4, 2000, pages 245-248, XP000996426 see Table 1	62-64
P,X	WO 00 62778 A (BRISTOL MYERS SQUIBB CO) 26 October 2000 (2000-10-26) see example 373	58
X	WO 99 24404 A (AMGEN INC) 20 May 1999 (1999-05-20) see general formula and examples	66,67,71

INTERNATIONAL SEARCH REPORT

international application No.
PCT/US 01/02478

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 1-57 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound
2. ☒ Claims Nos.: 1-8, 10-16, 20, 24, 28, 42-57, 66, 66
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-8,10-16,20,24,28,42-57,66,66

Present claims 1-8,10-16,20,24,28,42-57,62 and 66 relate to an extremely large number of possible compounds. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds and methods claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely those parts relating to the compounds and methods of claims 9,17-19,21-23,25-27 and 29-41 and compounds of claims 58-61,63 and 67.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/US 01/02478

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9525723	A	28-09-1995	AT 168099 T AU 688473 B AU 1898195 A BR 9507105 A CN 1143954 A CZ 9602690 A DE 69503365 D DE 69503365 T EP 0750611 A HU 74778 A,B JP 9510471 T PL 316289 A US 5756524 A ZA 9502205 A	15-07-1998 12-03-1998 09-10-1995 09-09-1997 26-02-1997 11-12-1996 13-08-1998 07-01-1999 02-01-1997 28-02-1997 21-10-1997 06-01-1997 26-05-1998 31-10-1995
WO 9936391	A	22-07-1999	AU 2111199 A US 6191170 B	02-08-1999 20-02-2001
US 6028111	A	22-02-2000	US 6100299 A AU 723706 B AU 1988497 A AU 724909 B AU 2319697 A CA 2248109 A CA 2248125 A EP 1009402 A EP 0927030 A JP 2000507223 T JP 2000506169 T WO 9732582 A WO 9732576 A ZA 9701891 A ZA 9701892 A	08-08-2000 07-09-2000 22-09-1997 05-10-2000 22-09-1997 12-09-1997 12-09-1997 21-06-2000 07-07-1999 13-06-2000 23-05-2000 12-09-1997 12-09-1997 07-09-1998 07-09-1998
WO 0055114	A	21-09-2000	AU 3876600 A	04-10-2000
WO 0121598	A	29-03-2001	NONE	
US 4261730	A	14-04-1981	US 4347075 A	31-08-1982
GB 1573576	A	28-08-1980	AR 220111 A AU 516014 B AU 3187777 A BE 862113 A CA 1112645 A CH 632997 A CS 196411 B DD 133884 A DE 2757111 A DK 571377 A FR 2387960 A FR 2381029 A HU 179545 B IL 53671 A IT 1090352 B JP 53079871 A JP 61034421 B MY 16982 A	15-10-1980 14-05-1981 28-06-1979 21-06-1978 17-11-1981 15-11-1982 31-03-1980 31-01-1979 06-07-1978 23-06-1978 17-11-1978 15-09-1978 29-11-1982 31-05-1982 26-06-1985 14-07-1978 07-08-1986 31-12-1982